

## Synphilin-1 rabbit pAb

Cat No.:ES3536

For research use only

## Overview

Product Name Synphilin-1 rabbit pAb

Host species Rabbit

**Applications** WB;IHC;IF;ELISA **Species Cross-Reactivity** Human;Mouse

**Recommended dilutions** Western Blot: 1/500 - 1/2000.

Immunohistochemistry: 1/100 - 1/300. ELISA: 1/10000. Not yet tested in other applications. The antiserum was produced against synthesized

Immunogen The antiserum was produced against synthesized

peptide derived from human Synphilin-1. AA

range:797-846

**Specificity** Synphilin-1 Polyclonal Antibody detects endogenous

levels of Synphilin-1 protein.

Formulation Liquid in PBS containing 50% glycerol, 0.5% BSA and

0.02% sodium azide.

**Storage** Store at -20°C. Avoid repeated freeze-thaw cycles.

Protein Name Synphilin-1 Gene Name SNCAIP

**Cellular localization** Cytoplasm . Detected in cytoplasmic inclusion

bodies, together with SNCA.

**Purification** The antibody was affinity-purified from rabbit

antiserum by affinity-chromatography using

epitope-specific immunogen.

Clonality Polyclonal
Concentration 1 mg/ml
Observed band 100kD
Human Gene ID 9627
Human Swiss-Prot Number Q9Y6H5

Alternative Names SNCAIP; Synphilin-1; Sph1;

Alpha-synuclein-interacting protein

**Background** This gene encodes a protein containing several

protein-protein interaction domains, including ankyrin-like repeats, a coiled-coil domain, and an ATP/GTP-binding motif. The encoded protein




+86-27-59760950 ELKbio@ELKbiotech.com


www.elkbiotech.com




interacts with alpha-synuclein in neuronal tissue and may play a role in the formation of cytoplasmic inclusions and neurodegeneration. A mutation in this gene has been associated with Parkinson's disease. Alternative splicing results in multiple transcript variants. [provided by RefSeq, Apr 2015],


Western Blot analysis of various cells using Synphilin-1 Polyclonal Antibody





Immunohistochemical analysis of paraffin-embedded Human lung cancer. Antibody was diluted at 1:100(4° overnight). High-pressure and temperature Tris-EDTA,pH8.0 was used for antigen retrieval. Negetive contrl (right) obtaned from antibody was pre-absorbed by immunogen peptide.






+86-27-59760950









+86-27-59760950

ELKbio@ELKbiotech.com