

UDG rabbit pAb

Cat No.: ES3673

For research use only

Overview

Product Name UDG rabbit pAb

Host species Rabbit

Applications WB;IHC;IF;ELISA Species Cross-Reactivity Human;Mouse;Rat

Recommended dilutions Western Blot: 1/500 - 1/2000.

Immunohistochemistry: 1/100 - 1/300. ELISA: 1/40000. Not yet tested in other applications.

Immunogen The antiserum was produced against synthesized

peptide derived from human UNG. AA

range:191-240

Specificity UDG Polyclonal Antibody detects endogenous levels

of UDG protein.

Formulation Liquid in PBS containing 50% glycerol, 0.5% BSA and

0.02% sodium azide.

Storage Store at -20°C. Avoid repeated freeze-thaw cycles.

Protein Name Uracil-DNA glycosylase

Gene Name UNG

Cellular localization [Isoform 1]: Mitochondrion.; [Isoform 2]: Nucleus. **Purification** The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using

epitope-specific immunogen.

Clonality Polyclonal
Concentration 1 mg/ml
Observed band 35kD
Human Gene ID 7374
Human Swiss-Prot Number P13051

Alternative Names UNG; DGU; UNG1; UNG15; Uracil-DNA glycosylase;

UDG

Background This gene encodes one of several uracil-DNA

glycosylases. One important function of uracil-DNA

glycosylases is to prevent mutagenesis by

eliminating uracil from DNA molecules by cleaving


the N-glycosylic bond and initiating the




+86-27-59760950 ELKbio@ELKbiotech.com v



base-excision repair (BER) pathway. Uracil bases occur from cytosine deamination or misincorporation of dUMP residues. Alternative promoter usage and splicing of this gene leads to two different isoforms: the mitochondrial UNG1 and the nuclear UNG2. The UNG2 term was used as a previous symbol for the CCNO gene (GeneID 10309), which has been confused with this gene, in the literature and some databases. [provided by RefSeq, Nov 2010],



Western Blot analysis of various cells using UDG Polyclonal Antibody. Secondary antibody(catalog#:RS0002) was diluted at 1:20000




+86-27-59760950


Western Blot analysis of COLO205 cells using UDG Polyclonal Antibody. Secondary antibody(catalog#:RS0002) was diluted at 1:20000







Western blot analysis of lysates from HepG2 and COLO cells, using UNG Antibody. The lane on the right is blocked with the synthesized peptide.



Western blot analysis of the lysates from HepG2 cells using UNG antibody.

