

## ALDH3A1 rabbit pAb

Cat No.: ES3755

For research use only

## Overview

**Product Name** ALDH3A1 rabbit pAb

**Host species** Rabbit WB;ELISA **Applications Species Cross-Reactivity** Human;Rat

**Recommended dilutions** Western Blot: 1/500 - 1/2000. ELISA: 1/20000. Not

yet tested in other applications.

The antiserum was produced against synthesized **Immunogen** 

peptide derived from human ALDH3A1. AA

range:236-285

**Specificity** ALDH3A1 Polyclonal Antibody detects endogenous

levels of ALDH3A1 protein.

**Formulation** Liquid in PBS containing 50% glycerol, 0.5% BSA and

0.02% sodium azide.

Store at -20°C. Avoid repeated freeze-thaw cycles. Storage **Protein Name** Aldehyde dehydrogenase dimeric NADP-preferring

**Gene Name** ALDH3A1 **Cellular localization** Cytoplasm.

Purification The antibody was affinity-purified from rabbit

antiserum by affinity-chromatography using

epitope-specific immunogen.

Clonality Polyclonal Concentration 1 mg/ml **Observed band** 50kD **Human Gene ID** 218 **Human Swiss-Prot Number** P30838

**Alternative Names** ALDH3A1; ALDH3; Aldehyde dehydrogenase, dimeric

> NADP-preferring; ALDHIII; Aldehyde dehydrogenase 3; Aldehyde dehydrogenase family 3 member A1 Aldehyde dehydrogenases oxidize various aldehydes

> to the corresponding acids. They are involved in the detoxification of alcohol-derived acetaldehyde and in the metabolism of corticosteroids, biogenic

amines, neurotransmitters, and lipid peroxidation.




**Background** 

+86-27-59760950 ELKbio@ELKbiotech.com



The enzyme encoded by this gene forms a cytoplasmic homodimer that preferentially oxidizes aromatic and medium-chain (6 carbons or more) saturated and unsaturated aldehyde substrates. It is thought to promote resistance to UV and 4-hydroxy-2-nonenal-induced oxidative damage in the cornea. The gene is located within the Smith-Magenis syndrome region on chromosome 17. Multiple alternatively spliced variants, encoding the same protein, have been identified. [provided by RefSeq, Sep 2008],



Western Blot analysis of extracts from A549, K562 cells, using ALDH3A1 Polyclonal Antibody. Secondary antibody(catalog#:RS0002) was diluted at 1:20000

