

## ABCG1 rabbit pAb

Cat No.: ES3790

For research use only

## Overview

Product Name ABCG1 rabbit pAb

Host species Rabbit

Applications WB;IHC;IF;ELISA Species Cross-Reactivity Human;Mouse;Rat

**Recommended dilutions** Western Blot: 1/500 - 1/2000. IHC-p: 1:100-300

ELISA: 1/10000. Not yet tested in other applications.

**Immunogen** Synthesized peptide derived from ABCG1 . at AA

range: 560-640

**Specificity** ABCG1 Polyclonal Antibody detects endogenous

levels of ABCG1 protein.

Formulation Liquid in PBS containing 50% glycerol, 0.5% BSA and

0.02% sodium azide.

Storage Store at -20°C. Avoid repeated freeze-thaw cycles.

Protein Name ATP-binding cassette sub-family G member 1

Gene Name ABCG1

**Cellular localization** Endoplasmic reticulum membrane ; Multi-pass

membrane protein . Golgi apparatus membrane ; Multi-pass membrane protein . Cell membrane . Predominantly localized in the intracellular compartments mainly associated with the

endoplasmic reticulum (ER) and Gol

**Purification** The antibody was affinity-purified from rabbit

antiserum by affinity-chromatography using

epitope-specific immunogen.

Clonality Polyclonal
Concentration 1 mg/ml
Observed band 75kD
Human Gene ID 9619
Human Swiss-Prot Number P45844

+86-27-59760950

Alternative Names ABCG1; ABC8; WHT1; ATP-binding cassette

sub-family G member 1; ATP-binding cassette

transporter 8; White protein homolog

**Background** The protein encoded by this gene is a member of

ELKbio@ELKbiotech.com




23-2, No.388 Gaoxin 2nd Road, Wuhan East Lake Hi-tech Development Zone, Hubei , P.R.



the superfamily of ATP-binding cassette (ABC) transporters. ABC proteins transport various molecules across extra- and intra-cellular membranes. ABC genes are divided into seven distinct subfamilies (ABC1, MDR/TAP, MRP, ALD, OABP, GCN20, White). This protein is a member of the White subfamily. It is involved in macrophage cholesterol and phospholipids transport, and may regulate cellular lipid homeostasis in other cell types. Six alternative splice variants have been identified. [provided by RefSeq, Jul 2008],

Western Blot analysis of extracts from K562 cells, using ABCG1 Polyclonal Antibody. Secondary antibody(catalog#:RS0002) was diluted at 1:20000



Immunohistochemical analysis of paraffin-embedded rat-brain, antibody was diluted at 1:100







Immunohistochemical analysis of paraffin-embedded rat-brain, antibody was diluted at 1:100



+86-27-59760950