

## UHRF1 rabbit pAb

## Cat No.:ES3796

For research use only

## Overview

Product Name	UHRF1 rabbit pAb
Host species	Rabbit
Applications	WB;ELISA
Species Cross-Reactivity	Human;Rat;Mouse;
<b>Recommended dilutions</b>	Western Blot: 1/500 - 1/2000. ELISA: 1/40000. Not
	yet tested in other applications.
Immunogen	Synthesized peptide derived from the Internal
_	region of human UHRF1.
Specificity	UHRF1 Polyclonal Antibody detects endogenous
	levels of UHRF1 protein.
Formulation	Liquid in PBS containing 50% glycerol, 0.5% BSA and
	0.02% sodium azide.
Storage	Store at -20°C. Avoid repeated freeze-thaw cycles.
Protein Name	E3 ubiquitin-protein ligase UHRF1
Gene Name	UHRF1
Cellular localization	Nucleus . Localizes to replication foci. Enriched in
	pericentric heterochromatin. Also localizes to
	euchromatic regions.
Purification	The antibody was affinity-purified from rabbit
	antiserum by affinity-chromatography using
	epitope-specific immunogen.
Clonality	Polyclonal
Concentration	1 mg/ml
Observed band	89kD
Human Gene ID	29128
Human Swiss-Prot Number	Q96T88
Alternative Names	UHRF1; ICBP90; NP95; RNF106; E3 ubiquitin-protein
	ligase UHRF1; Inverted CCAAT box-binding protein of
	90 kDa; Nuclear protein 95; Nuclear zinc finger
	protein Np95; HuNp95; hNp95; RING finger protein
	106;Transcription factor ICBP90; Ubiquitin-like PHD
	and
Background	This gene encodes a member of a subfamily of
-	



+86-27-59760950


ELKbio@ELKbiotech.com

www.elkbiotech.com

23-2, No.388 Gaoxin 2nd Road, Wuhan East Lake Hi-tech Development Zone, Hubei , P.R.C



RING-finger type E3 ubiquitin ligases. The protein binds to specific DNA sequences, and recruits a histone deacetylase to regulate gene expression. Its expression peaks at late G1 phase and continues during G2 and M phases of the cell cycle. It plays a major role in the G1/S transition by regulating topoisomerase IIalpha and retinoblastoma gene expression, and functions in the p53-dependent DNA damage checkpoint. It is regarded as a hub protein for the integration of epigenetic information. This gene is up-regulated in various cancers, and it is therefore considered to be a therapeutic target. Multiple transcript variants encoding different isoforms have been found for this gene. A related pseudogene exists on chromosome 12. [provided by RefSeq, Feb 2014],



Western Blot analysis of extracts from Jurkat cells, using UHRF1 Polyclonal Antibody. Secondary antibody(catalog#:RS0002) was diluted at 1:20000 cells nucleus extracted by Minute TM Cytoplasmic and Nuclear Fractionation kit (SC-003,Inventbiotech,MN,USA).



+86-27-59760950

ELKbio@ELKbiotech.com

www.elkbiotech.com

23-2, No.388 Gaoxin 2nd Road, Wuhan East Lake Hi-tech Development Zone, Hubei , P.R.C