

ADM rabbit pAb

Cat No.: ES3932

For research use only

Overview

Product Name ADM rabbit pAb

Host species Rabbit

Applications WB;IHC;IF;ELISA **Species Cross-Reactivity** Human;Rat

Recommended dilutions Western Blot: 1/500 - 1/2000. IHC-p: 1:100-300

ELISA: 1/20000. Not yet tested in other applications.

The antiserum was produced against synthesized **Immunogen**

peptide derived from the Internal region of human

ADM. AA range:101-150

Specificity ADM Polyclonal Antibody detects endogenous levels

of ADM protein.

Formulation Liquid in PBS containing 50% glycerol, 0.5% BSA and

0.02% sodium azide.

Storage Store at -20°C. Avoid repeated freeze-thaw cycles.

Protein Name ADM **Gene Name** ADM **Cellular localization** Secreted.

Purification The antibody was affinity-purified from rabbit

antiserum by affinity-chromatography using

epitope-specific immunogen.

Clonality Polyclonal Concentration 1 mg/ml **Observed band** 20kD **Human Gene ID** 133 **Human Swiss-Prot Number** P35318

Alternative Names ADM; AM; ADM

Background The protein encoded by this gene is a

> preprohormone which is cleaved to form two biologically active peptides, adrenomedullin and proadrenomedullin N-terminal 20 peptide. Adrenomedullin is a 52 aa peptide with several


functions, including vasodilation, regulation of hormone secretion, promotion of angiogenesis, and




+86-27-59760950 ELKbio@ELKbiotech.com




antimicrobial activity. The antimicrobial activity is antibacterial, as the peptide has been shown to kill E. coli and S. aureus at low concentration. [provided by RefSeq, Aug 2014],



Western Blot analysis of mouse kidney, mouse lung, mouse heart cells using ADM Polyclonal Antibody. Secondary antibody(catalog#:RS0002) was diluted at 1:20000



Immunohistochemical analysis of paraffin-embedded human-liver, antibody was diluted at 1:100




+86-27-59760950

Immunohistochemical analysis of paraffin-embedded human-liver, antibody was diluted at 1:100







Immunohistochemical analysis of paraffin-embedded rat-kidney, antibody was diluted at 1:100



+86-27-59760950