

DPYD rabbit pAb

Cat No.:ES3957

For research use only

Overview

Product Name	DPYD rabbit pAb
Host species	Rabbit
Applications	WB;IHC;IF;ELISA
Species Cross-Reactivity	Human;Mouse;Rat
Recommended dilutions	Western Blot: 1/500 - 1/2000. IHC-p: 1:100-300
	ELISA: 1/20000. Not yet tested in other applications.
Immunogen	The antiserum was produced against synthesized
2	peptide derived from the Internal region of human
	DPYD. AA range:351-400
Specificity	DPYD Polyclonal Antibody detects endogenous
	levels of DPYD protein.
Formulation	Liquid in PBS containing 50% glycerol, 0.5% BSA and
	0.02% sodium azide.
Storage	Store at -20°C. Avoid repeated freeze-thaw cycles.
Protein Name	Dihydropyrimidine dehydrogenase [NADP(+)]
Gene Name	DPYD
Cellular localization	Cytoplasm.
Purification	The antibody was affinity-purified from rabbit
	antiserum by affinity-chromatography using
	epitope-specific immunogen.
Clonality	Polyclonal
Concentration	1 mg/ml
Observed band	120kD
Human Gene ID	1806
Human Swiss-Prot Number	Q12882
Alternative Names	DPYD; Dihydropyrimidine dehydrogenase [NADP(+)];
	DHPDHase; DPD; Dihydrothymine dehydrogenase;
	Dihydrouracil dehydrogenase
Background	The protein encoded by this gene is a pyrimidine
	catabolic enzyme and the initial and rate-limiting
	factor in the pathway of uracil and thymidine
	catabolism. Mutations in this gene result in
	dihydropyrimidine dehydrogenase deficiency, an



+86-27-59760950

ELKbio@ELKbiotech.com


www.elkbiotech.com


23-2, No.388 Gaoxin 2nd Road, Wuhan East Lake Hi-tech Development Zone, Hubei , P.R.C



error in pyrimidine metabolism associated with thymine-uraciluria and an increased risk of toxicity in cancer patients receiving 5-fluorouracil chemotherapy. Two transcript variants encoding different isoforms have been found for this gene. [provided by RefSeq, May 2009],


Western Blot analysis of HepG2 cells using DPYD Polyclonal Antibody. Secondary antibody(catalog#:RS0002) was diluted at 1:20000







Immunohistochemical analysis of paraffin-embedded mouse-lung, antibody was diluted at 1:100





+86-27-59760950


ELKbio@ELKbiotech.com

www.elkbiotech.com

23-2, No.388 Gaoxin 2nd Road, Wuhan East Lake Hi-tech Development Zone, Hubei , P.R.C



Immunohistochemical analysis of paraffin-embedded mouse-lung, antibody was diluted at 1:100





+86-27-59760950

ELKbio@ELKbiotech.com

www.elkbiotech.com

23-2, No.388 Gaoxin 2nd Road, Wuhan East Lake Hi-tech Development Zone, Hubei , P.R.C