

PSP rabbit pAb

Cat No.: ES4025

For research use only

Overview

Product Name PSP rabbit pAb

Host species Rabbit

Applications WB;ELISA;IHC

Species Cross-Reactivity Human;Rat;Mouse;

Recommended dilutions WB 1:500-2000;IHC-p 1:50-300; ELISA 2000-20000 **Immunogen** The antiserum was produced against synthesized

peptide derived from the C-terminal region of

human PSPN. AA range:101-150

Specificity PSP Polyclonal Antibody detects endogenous levels

of PSP protein.

Formulation Liquid in PBS containing 50% glycerol, 0.5% BSA and

0.02% sodium azide.

Storage Store at -20°C. Avoid repeated freeze-thaw cycles.

Protein Name Persephin
Gene Name PSPN
Cellular localization Secreted.

Purification The antibody was affinity-purified from rabbit

antiserum by affinity-chromatography using

epitope-specific immunogen.

Clonality Polyclonal
Concentration 1 mg/ml
Observed band 16kD
Human Gene ID 5623
Human Swiss-Prot Number O60542

Alternative Names PSPN; Persephin; PSP

Background persephin(PSPN) Homo sapiens This gene

encodes a secreted ligand of the GDNF (glial cell line-derived neurotrophic factor) subfamily and TGF-beta (transforming growth factor-beta)

superfamily of proteins. The encoded preproprotein is proteolytically processed to generate the mature

protein. This protein signals through the RET receptor tyrosine kinase and a GPI-linked




+86-27-59760950 ELKbio@ELKbiotech.com

www.elkbiotech.com




coreceptor, and promotes survival of neuronal populations. This protein may play a role in cell death, and nervous system development and function. Elevated expression of this gene has been observed in oral squamous cell carcinoma. [provided by RefSeq, Aug 2016],


Western Blot analysis of 293 cells using PSP Polyclonal Antibody. Secondary antibody(catalog#:RS0002) was diluted at 1:20000



Western blot analysis of lysate from 293 cells, using PSPN Antibody.



Immunohistochemical analysis of paraffin-embedded human liver cancer. 1, Antibody was diluted at 1:200(4° overnight). 2, Tris-EDTA,pH9.0 was used for antigen retrieval. 3,Secondary antibody was diluted at 1:200(room temperature, 45min).



+86-27-59760950