

CD81 rabbit pAb

Cat No.:ES4124

For research use only

Overview

Product Name	CD81 rabbit pAb
Host species	Rabbit
Applications	IF;WB;ELISA
Species Cross-Reactivity	Human;Mouse;Rat
Recommended dilutions	IF: 1:50-200 Western Blot: 1/500 - 1/2000. ELISA:
	1/20000. Not yet tested in other applications.
Immunogen	The antiserum was produced against synthesized
	peptide derived from the Internal region of human
	CD81. AA range:111-160
Specificity	CD81 Polyclonal Antibody detects endogenous levels
	of CD81 protein.
Formulation	Liquid in PBS containing 50% glycerol, 0.5% BSA and
	0.02% sodium azide.
Storage	Store at -20 $^\circ\!\mathrm{C}$. Avoid repeated freeze-thaw cycles.
Protein Name	CD81 antigen
Gene Name	CD81
Cellular localization	Cell membrane ; Multi-pass membrane protein .
	Basolateral cell membrane ; Multi-pass membrane
	protein . Associates with CLDN1 and the
	CLDN1-CD81 complex localizes to the basolateral cell
	membrane
Purification	The antibody was affinity-purified from rabbit
	antiserum by affinity-chromatography using
	epitope-specific immunogen.
Clonality	Polyclonal
Concentration	1 mg/ml
Observed band	26kD
Human Gene ID	975
Human Swiss-Prot Number	P60033
Alternative Names	CD81; TAPA1; TSPAN28; CD81 antigen; 26 kDa cell
	surface protein TAPA-1; Target of the
	antiproliferative antibody 1; Tetraspanin-28;
	Tspan-28; CD81



+86-27-59760950

ELKbio@ELKbiotech.com

www.elkbiotech.com

23-2, No.388 Gaoxin 2nd Road, Wuhan East Lake Hi-tech Development Zone, Hubei , P.R.C



Background

The protein encoded by this gene is a member of the transmembrane 4 superfamily, also known as the tetraspanin family. Most of these members are cell-surface proteins that are characterized by the presence of four hydrophobic domains. The proteins mediate signal transduction events that play a role in the regulation of cell development, activation, growth and motility. This encoded protein is a cell surface glycoprotein that is known to complex with integrins. This protein appears to promote muscle cell fusion and support myotube maintenance. Also it may be involved in signal transduction. This gene is localized in the tumor-suppressor gene region and thus it is a candidate gene for malignancies. Two transcript variants encoding different isoforms have been found for this gene. [provided by RefSeq, Jul 2014],



+86-27-59760950

ELKbio@ELKbiotech.com

www.elkbiotech.com

23-2, No.388 Gaoxin 2nd Road, Wuhan East Lake Hi-tech Development Zone, Hubei , P.R.C