

Integrin αL rabbit pAb

Cat No.: ES4308

For research use only

Overview

Product Name Integrin αL rabbit pAb

Host species Rabbit
Applications WB;ELISA

Species Cross-Reactivity Human;Rat;Mouse;

Recommended dilutions Western Blot: 1/500 - 1/2000. ELISA: 1/10000. Not

yet tested in other applications.

Immunogen The antiserum was produced against synthesized

peptide derived from the Internal region of human

ITGAL. AA range:171-220

Specificity Integrin αL Polyclonal Antibody detects endogenous

levels of Integrin αL protein.

Formulation Liquid in PBS containing 50% glycerol, 0.5% BSA and

0.02% sodium azide.

Storage Store at -20°C. Avoid repeated freeze-thaw cycles.

Protein Name Integrin alpha-L

Gene Name ITGAL

Cell ular localization Cell membrane ; Single-pass type I membrane

protein.

Purification The antibody was affinity-purified from rabbit

antiserum by affinity-chromatography using

epitope-specific immunogen.

Clonality Polyclonal
Concentration 1 mg/ml
Observed band 130kD
Human Gene ID 3683
Human Swiss-Prot Number P20701

Alternative Names ITGAL; CD11A; Integrin alpha-L; CD11 antigen-like

family member A; Leukocyte adhesion glycoprotein

LFA-1 alpha chain; LFA-1A; Leukocyte

function-associated molecule 1 alpha chain; CD11a

Background integrin subunit alpha L(ITGAL) Homo sapiens

ITGAL encodes the integrin alpha L chain. Integrins

are heterodimeric integral membrane proteins




+86-27-59760950 ELKbio@ELKbiotech.com www.elkbiotech.co



I-domain containing alpha integrin combines with the beta 2 chain (ITGB2) to form the integrin lymphocyte function-associated antigen-1 (LFA-1), which is expressed on all leukocytes. LFA-1 plays a central role in leukocyte intercellular adhesion through interactions with its ligands, ICAMs 1-3 (intercellular adhesion molecules 1 through 3), and also functions in lymphocyte costimulatory signaling. Two transcript variants encoding different isoforms have been found for this gene. [provided by RefSeq, Jul 2008],

composed of an alpha chain and a beta chain. This

Western Blot analysis of 293 cells using Integrin αL Polyclonal Antibody. Secondary antibody(catalog#:RS0002) was diluted at 1:20000



+86-27-59760950

