

T-cadherin rabbit pAb

Cat No.: ES4425

For research use only

Overview

Product Name T-cadherin rabbit pAb

Host species Rabbit

Applications WB;IHC;IF;ELISA Species Cross-Reactivity Human;Mouse;Rat

Recommended dilutions Western Blot: 1/500 - 1/2000.

Immunohistochemistry: 1/100 - 1/300. Immunofluorescence: 1/200 - 1/1000. ELISA: 1/20000. Not yet tested in other applications.

Immunogen The antiserum was produced against synthesized

peptide derived from human CDH13. AA

range:331-380

Specificity T-cadherin Polyclonal Antibody detects endogenous

levels of T-cadherin protein.

Formulation Liquid in PBS containing 50% glycerol, 0.5% BSA and

0.02% sodium azide.

Storage Store at -20°C. Avoid repeated freeze-thaw cycles.

Protein Name Cadherin-13
Gene Name CDH13

Cellular localizationCell membrane; Lipid-anchor, GPI-anchor.PurificationThe antibody was affinity-purified from rabbitanticorum by affinity shromatography using

antiserum by affinity-chromatography using

epitope-specific immunogen.

Clonality Polyclonal
Concentration 1 mg/ml
Observed band 78kD
Human Gene ID 1012
Human Swiss-Prot Number P55290

Alternative Names CDH13; CDHH; Cadherin-13; Heart cadherin;

H-cadherin; P105; Truncated cadherin; T-cad;

T-cadherin

Background This gene encodes a member of the cadherin

superfamily. The encoded protein is localized to the surface of the cell membrane and is anchored by a






GPI moiety, rather than by a transmembrane domain. The protein lacks the cytoplasmic domain characteristic of other cadherins, and so is not thought to be a cell-cell adhesion glycoprotein. This protein acts as a negative regulator of axon growth during neural differentiation. It also protects vascular endothelial cells from apoptosis due to oxidative stress, and is associated with resistance to atherosclerosis. The gene is hypermethylated in many types of cancer. Alternative splicing results in multiple transcript variants encoding different isoforms. [provided by RefSeq, May 2011],



40-35--25-

15--

Western Blot analysis of LOVO cells using T-cadherin Polyclonal Antibody




+86-27-59760950

Immunohistochemistry analysis of paraffin-embedded human heart tissue, using CDH13 Antibody. The picture on the right is blocked with the synthesized peptide.







Western blot analysis of lysates from LOVO cells, using CDH13 Antibody. The lane on the right is blocked with the synthesized peptide.



+86-27-59760950