



# ADAMTS-7 rabbit pAb

Cat No.:ES4607

For research use only

## Overview

<b>Product Name</b>	ADAMTS-7 rabbit pAb
<b>Host species</b>	Rabbit
<b>Applications</b>	IHC;IF;ELISA
<b>Species Cross-Reactivity</b>	Human;Rat;Mouse;
<b>Recommended dilutions</b>	Immunohistochemistry: 1/100 - 1/300. ELISA: 1/40000. Not yet tested in other applications.
<b>Immunogen</b>	Synthesized peptide derived from ADAMTS-7 . at AA range: 150-230
<b>Specificity</b>	ADAMTS-7 Polyclonal Antibody detects endogenous levels of ADAMTS-7 protein.
<b>Formulation</b>	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.
<b>Storage</b>	Store at -20°C. Avoid repeated freeze-thaw cycles.
<b>Protein Name</b>	A disintegrin and metalloproteinase with thrombospondin motifs 7
<b>Gene Name</b>	ADAMTS7
<b>Cellular localization</b>	Secreted, extracellular space, extracellular matrix . Also found associated with the external cell surface. .
<b>Purification</b>	The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen.
<b>Clonality</b>	Polyclonal
<b>Concentration</b>	1 mg/ml
<b>Observed band</b>	
<b>Human Gene ID</b>	11173
<b>Human Swiss-Prot Number</b>	Q9UKP4
<b>Alternative Names</b>	ADAMTS7; A disintegrin and metalloproteinase with thrombospondin motifs 7; ADAM-TS 7; ADAM-TS7; ADAMTS-7; COMPase
<b>Background</b>	The protein encoded by this gene is a member of the ADAMTS (a disintegrin and metalloproteinase with thrombospondin motifs) family. Members of





this family share several distinct protein modules, including a propeptide region, a metalloproteinase domain, a disintegrin-like domain, and a thrombospondin type 1 (TS) motif. Individual members of this family differ in the number of C-terminal TS motifs, and some have unique C-terminal domains. The encoded preproprotein is proteolytically processed to generate the mature enzyme. This enzyme contains two C-terminal TS motifs and may regulate vascular smooth muscle cell (VSMC) migration. Mutations in this gene may be associated with susceptibility to coronary artery disease. [provided by RefSeq, Feb 2016],

Immunohistochemical analysis of paraffin-embedded human tonsil. 1, Antibody was diluted at 1:200(4° overnight). 2, Tris-EDTA,pH9.0 was used for antigen retrieval. 3,Secondary antibody was diluted at 1:200(room temperature, 30min).

