

## Olfactory receptor 4L1 rabbit pAb

Cat No.: ES4703

For research use only

## Overview

Product Name Olfactory receptor 4L1 rabbit pAb

Host species Rabbit
Applications WB;IF;ELISA

**Species Cross-Reactivity** Human; Rat; Mouse;

**Recommended dilutions** Western Blot: 1/500 - 1/2000. Immunofluorescence:

1/200 - 1/1000. ELISA: 1/10000. Not yet tested in

other applications.

Immunogen The antiserum was produced against synthesized

peptide derived from human OR4L1. AA

range:262-311

**Specificity** Olfactory receptor 4L1 Polyclonal Antibody detects

endogenous levels of Olfactory receptor 4L1 protein.

Formulation Liquid in PBS containing 50% glycerol, 0.5% BSA and

0.02% sodium azide.

**Storage** Store at -20°C. Avoid repeated freeze-thaw cycles.

Protein Name Olfactory receptor 4L1

Gene Name OR4L1

Cellular localizationCell membrane; Multi-pass membrane protein.PurificationThe antibody was affinity-purified from rabbit

antiserum by affinity-chromatography using

epitope-specific immunogen.

ClonalityPolyclonalConcentration1 mg/mlObserved band34kDHuman Gene ID122742Human Swiss-Prot NumberQ8NH43

Alternative Names OR4L1; OR4L2P; Olfactory receptor 4L1; Olfactory

receptor 4L2; Olfactory receptor OR14-28

**Background** olfactory receptor family 4 subfamily L member

1(OR4L1) Homo sapiens Olfactory receptors interact with odorant molecules in the nose, to initiate a neuronal response that triggers the perception of a smell. The olfactory receptor




+86-27-59760950 ELKbio@ELKbiotech.com


www.elkbiotech.com




proteins are members of a large family of G-protein-coupled receptors (GPCR) arising from single coding-exon genes. Olfactory receptors share a 7-transmembrane domain structure with many neurotransmitter and hormone receptors and are responsible for the recognition and G protein-mediated transduction of odorant signals. The olfactory receptor gene family is the largest in the genome. The nomenclature assigned to the olfactory receptor genes and proteins for this organism is independent of other organisms. [provided by RefSeq, Jul 2008],



Western Blot analysis of K562 cells using Olfactory receptor 4L1 Polyclonal Antibody







+86-27-59760950







Western blot analysis of lysates from K562 and MCF-7 cells, using OR4L1 Antibody. The lane on the right is blocked with the synthesized peptide.



Western blot analysis of the lysates from HepG2 cells using OR4L1 antibody.

