

## Claudin-7 rabbit pAb

Cat No.:ES4794

For research use only

## Overview

**Product Name** Claudin-7 rabbit pAb

**Host species** Rabbit

WB;IHC;IF;ELISA **Applications Species Cross-Reactivity** Human; Mouse; Rat

**Recommended dilutions** Western Blot: 1/500 - 1/2000.

> Immunohistochemistry: 1/100 - 1/300. ELISA: 1/10000. Not yet tested in other applications.

The antiserum was produced against synthesized **Immunogen** 

peptide derived from human Claudin 7. AA

range:162-211

Specificity Claudin-7 Polyclonal Antibody detects endogenous

levels of Claudin-7 protein.

**Formulation** Liquid in PBS containing 50% glycerol, 0.5% BSA and

0.02% sodium azide.

Store at -20°C. Avoid repeated freeze-thaw cycles. **Storage** 

**Protein Name** Claudin-7 **Gene Name** CLDN7

Cellular localization Cell membrane; Multi-pass membrane protein.

> Basolateral cell membrane . Cell junction, tight junction. Co-localizes with EPCAM at the basolateral

cell membrane and tight junction. .

Purification The antibody was affinity-purified from rabbit

antiserum by affinity-chromatography using

epitope-specific immunogen.

Polyclonal Clonality Concentration 1 mg/ml **Observed band** 32kD **Human Gene ID** 1366 **Human Swiss-Prot Number** O95471

**Alternative Names** CLDN7; CEPTRL2; CPETRL2; Claudin-7; CLDN-7 **Background** This gene encodes a member of the claudin family.

Claudins are integral membrane proteins and

components of tight junction strands. Tight junction




+86-27-59760950 ELKbio@ELKbiotech.com



strands serve as a physical barrier to prevent solutes and water from passing freely through the paracellular space between epithelial or endothelial cell sheets, and also play critical roles in maintaining cell polarity and signal transductions. Differential expression of this gene has been observed in different types of malignancies, including breast cancer, ovarian cancer, hepatocellular carcinomas, urinary tumors, prostate cancer, lung cancer, head and neck cancers, thyroid carcinomas, etc.. Alternatively spliced transcript variants encoding different isoforms have been found.[provided by RefSeq, May 2010],

Immunohistochemistry analysis of paraffin-embedded human prostate carcinoma tissue, using Claudin 7 Antibody. The picture on the right is blocked with the synthesized peptide.



-- 117 -- 85 -- 48 -- 34 -- 26 -- 19 (kD)

+86-27-59760950

Western blot analysis of lysates from rat liver cells, using Claudin 7 Antibody. The lane on the right is blocked with the synthesized peptide.

