

DGAT2L3 rabbit pAb

Cat No.: ES4950

For research use only

Overview

Product Name DGAT2L3 rabbit pAb

Host species Rabbit
Applications WB;ELISA

Species Cross-Reactivity Human;Rat;Mouse;

Recommended dilutions Western Blot: 1/500 - 1/2000. ELISA: 1/5000. Not

yet tested in other applications.

Immunogen The antiserum was produced against synthesized

peptide derived from human AWAT1. AA

range:261-310

Specificity DGAT2L3 Polyclonal Antibody detects endogenous

levels of DGAT2L3 protein.

Formulation Liquid in PBS containing 50% glycerol, 0.5% BSA and

0.02% sodium azide.

Storage Store at -20°C. Avoid repeated freeze-thaw cycles.

Protein Name Acyl-CoA wax alcohol acyltransferase 1

Gene Name AWAT1

Cellular localization Endoplasmic reticulum membrane ; Multi-pass

membrane protein.

Purification The antibody was affinity-purified from rabbit

antiserum by affinity-chromatography using

epitope-specific immunogen.

ClonalityPolyclonalConcentration1 mg/mlObserved band39kDHuman Gene ID158833Human Swiss-Prot NumberQ58HT5

Alternative Names AWAT1; DGA2; DGAT2L3; Acyl-CoA wax alcohol

acyltransferase 1; Diacylglycerol O-acyltransferase 2-like protein 3; Diacylglycerol acyltransferase 2; Long-chain-alcohol O-fatty-acyltransferase 1

Background The protein encoded by this gene belongs to the

diacylglycerol acyltransferase family. It esterifies long chain (wax) alcohols with acyl-CoA-derived




+86-27-59760950 ELKbio@ELKbiotech.com

www.elkbiotech.com




fatty acids to produce wax esters. Wax esters are enriched in sebum, suggesting that this enzyme plays a central role in lipid metabolism in skin. Consistent with this observation, this protein is predominantly expressed in the sebaceous gland of the skin. [provided by RefSeq, Sep 2009],


Western Blot analysis of 293 cells using DGAT2L3 Polyclonal Antibody diluted at 1:1000



Western Blot analysis of 293T NIH-3T3 COLO cells using DGAT2L3 Polyclonal Antibody diluted at 1:1000




Western blot analysis of mouse-kidney KB lysis using DGAT2L3 antibody. Antibody was diluted at 1:1000









Western blot analysis of lysates from 293 cells, using AWAT1 Antibody. The lane on the right is blocked with the synthesized peptide.

