

## MKP-4 rabbit pAb

## Cat No.:ES5033

For research use only

## Overview

Product Name	MKP-4 rabbit pAb	
Host species	Rabbit	
Applications	WB;IHC;IF;ELISA	
Species Cross-Reactivity	Human; Mouse; Rat	
Recommended dilutions	Western Blot: 1/500 - 1/2000.	
	Immunohistochemistry: 1/100 - 1/300.	
	Immunofluorescence: 1/200 - 1/1000. ELISA:	
	1/20000. Not yet tested in other applications.	
Immunogen	The antiserum was produced against synthesized	
	peptide derived from human DUSP9. AA	
	range:151-200	
Specificity	MKP-4 Polyclonal Antibody detects endogenous	
	levels of MKP-4 protein.	
Formulation	Liquid in PBS containing 50% glycerol, 0.5% BSA and	
	0.02% sodium azide.	
Storage	Store at -20 $^\circ\!\mathrm{C}$ . Avoid repeated freeze-thaw cycles.	
Protein Name	Dual specificity protein phosphatase 9	
Gene Name	DUSP9	
Cellular localization	Cytoplasm.	
Purification	The antibody was affinity-purified from rabbit	
	antiserum by affinity-chromatography using	
	epitope-specific immunogen.	
Clonality	Polyclonal	
Concentration	1 mg/ml	
Observed band	42kD	
Human Gene ID	1852	
Human Swiss-Prot Number	Q99956	
Alternative Names	DUSP9; MKP4; Dual specificity protein phosphatase	
	9; Mitogen-activated protein kinase phosphatase 4;	
	MAP kinase phosphatase 4; MKP-4	
Background	The protein encoded by this gene is a member of	
	the dual specificity protein phosphatase subfamily.	
	These phosphatases inactivate their target kinases	



+86-27-59760950

ELKbio@ELKbiotech.com

www.elkbiotech.com

23-2, No.388 Gaoxin 2nd Road, Wuhan East Lake Hi-tech Development Zone, Hubei , P.R.C



## by dephosphorylating both the

phosphoserine/threonine and phosphotyrosine residues. They negatively regulate members of the mitogen-activated protein (MAP) kinase superfamily (MAPK/ERK, SAPK/JNK, p38), which is associated with cellular proliferation and differentiation. Different members of the family of dual specificity phosphatases show distinct substrate specificities for various MAP kinases, different tissue distribution and subcellular localization, and different modes of inducibility of their expression by extracellular stimuli. This gene product shows selectivity for members of the ERK family of MAP kinases and is localized to the cytoplasm and nucleus. Aberrant expression of this gene is associated with type 2 diabetes and cancer progr



+86-27-59760950

ELKbio@ELKbiotech.com

www.elkbiotech.com

23-2, No.388 Gaoxin 2nd Road, Wuhan East Lake Hi-tech Development Zone, Hubei , P.R.C