

Olfactory receptor 4C16 rabbit pAb

Cat No.: ES5207

For research use only

Overview

Product Name Olfactory receptor 4C16 rabbit pAb

Host species Rabbit
Applications WB;IF;ELISA

Species Cross-Reactivity Human; Rat; Mouse;

Recommended dilutions Western Blot: 1/500 - 1/2000. Immunofluorescence:

1/200 - 1/1000. ELISA: 1/20000. Not yet tested in

other applications.

Immunogen The antiserum was produced against synthesized

peptide derived from human OR4C16. AA

range:261-310

Specificity Olfactory receptor 4C16 Polyclonal Antibody detects

endogenous levels of Olfactory receptor 4C16

protein.

Formulation Liquid in PBS containing 50% glycerol, 0.5% BSA and

0.02% sodium azide.

Storage Store at -20°C. Avoid repeated freeze-thaw cycles.

Protein Name Olfactory receptor 4C16

Gene Name OR4C16

Cellular localizationCell membrane; Multi-pass membrane protein.PurificationThe antibody was affinity-purified from rabbit
antiserum by affinity-chromatography using

epitope-specific immunogen.

Clonality Polyclonal
Concentration 1 mg/ml
Observed band 35kD
Human Gene ID 219428
Human Swiss-Prot Number O8NGL9

Alternative Names OR4C16; Olfactory receptor 4C16; Olfactory

receptor OR11-135

Background Olfactory receptors interact with odorant molecules


in the nose, to initiate a neuronal response that triggers the perception of a smell. The olfactory receptor proteins are members of a large family of




+86-27-59760950 ELKbio@ELKbiotech.com www.elkbiote



G-protein-coupled receptors (GPCR) arising from single coding-exon genes. Olfactory receptors share a 7-transmembrane domain structure with many neurotransmitter and hormone receptors and are responsible for the recognition and G protein-mediated transduction of odorant signals. The olfactory receptor gene family is the largest in the genome. The nomenclature assigned to the olfactory receptor genes and proteins for this organism is independent of other organisms. This olfactory receptor gene is a segregating pseudogene, where some individuals have an allele that encodes a functional olfactory receptor, while other individuals have an allele encoding a



Western Blot analysis of SH-SY5Y cells using Olfactory receptor 4C16 Polyclonal Antibody diluted at 1:500




+86-27-59760950

Immunofluorescence analysis of A549 cells, using OR4C16 Antibody. The picture on the right is blocked with the synthesized peptide.




23-2, No.388 Gaoxin 2nd Road, Wuhan East Lake Hi-tech Development Zone, Hubei, P.R.C





Western blot analysis of lysates from LOVO cells, using OR4C16 Antibody. The lane on the right is blocked with the synthesized peptide.



Western blot analysis of the lysates from K562 cells using OR4C16 antibody.

