

Olfactory receptor 8K3 rabbit pAb

Cat No.:ES5211

For research use only

Overview

Product Name	Olfactory receptor 8K3 rabbit pAb
Host species	Rabbit
Applications	WB;IF;ELISA
Species Cross-Reactivity	Human;Rat;Mouse;
Recommended dilutions	Western Blot: 1/500 - 1/2000. Immunofluorescence:
	1/200 - 1/1000. ELISA: 1/10000. Not yet tested in
	other applications.
Immunogen	The antiserum was produced against synthesized
	peptide derived from human OR8K3. AA
	range:236-285
Specificity	Olfactory receptor 8K3 Polyclonal Antibody detects
	endogenous levels of Olfactory receptor 8K3
	protein.
Formulation	Liquid in PBS containing 50% glycerol, 0.5% BSA and
	0.02% sodium azide.
Storage	Store at -20°C. Avoid repeated freeze-thaw cycles.
Protein Name	Olfactory receptor 8K3
Gene Name	OR8K3
Cellular localization	Cell membrane; Multi-pass membrane protein.
Purification	The antibody was affinity-purified from rabbit
	antiserum by affinity-chromatography using
	epitope-specific immunogen.
Clonality	Polyclonal
Concentration	1 mg/ml
Observed band	33kD
Human Gene ID	219473
Human Swiss-Prot Number	Q8NH51
Alternative Names	OR8K3; Olfactory receptor 8K3; Olfactory receptor
	OR11-181
Background	Olfactory receptors interact with odorant molecules
	in the nose, to initiate a neuronal response that
	triggers the perception of a smell. The olfactory
	receptor proteins are members of a large family of



+86-27-59760950

ELKbio@ELKbiotech.com


www.elkbiotech.com

23-2, No.388 Gaoxin 2nd Road, Wuhan East Lake Hi-tech Development Zone, Hubei , P.R.C




G-protein-coupled receptors (GPCR) arising from single coding-exon genes. Olfactory receptors share a 7-transmembrane domain structure with many neurotransmitter and hormone receptors and are responsible for the recognition and G protein-mediated transduction of odorant signals. The olfactory receptor gene family is the largest in the genome. The nomenclature assigned to the olfactory receptor genes and proteins for this organism is independent of other organisms. This olfactory receptor gene is a segregating pseudogene, where some individuals have an allele that encodes a functional olfactory receptor, while other individuals have an allele encoding a

Western Blot analysis of LOVO cells using Olfactory receptor 8K3 Polyclonal Antibody



Immunofluorescence analysis of HUVEC cells, using OR8K3 Antibody. The picture on the right is blocked with the synthesized peptide.






+86-27-59760950

ELKbio@ELKbiotech.com

www.elkbiotech.com

23-2, No.388 Gaoxin 2nd Road, Wuhan East Lake Hi-tech Development Zone, Hubei , P.R.G



Western blot analysis of lysates from LOVO cells, using OR8K3 Antibody. The lane on the right is blocked with the synthesized peptide.



+86-27-59760950

ELKbio@ELKbiotech.com

www.elkbiotech.com

23-2, No.388 Gaoxin 2nd Road, Wuhan East Lake Hi-tech Development Zone, Hubei , P.R.C