

## Olfactory receptor 4D6 rabbit pAb

Cat No.:ES5221

For research use only

## Overview

Product Name Olfactory receptor 4D6 rabbit pAb

Host species Rabbit
Applications WB;ELISA

**Species Cross-Reactivity** Human; Rat; Mouse;

Recommended dilutions Western Blot: 1/500 - 1/2000. ELISA: 1/20000. Not

yet tested in other applications.

Immunogen The antiserum was produced against synthesized

peptide derived from human OR4D6. AA

range:260-309

**Specificity** Olfactory receptor 4D6 Polyclonal Antibody detects

endogenous levels of Olfactory receptor 4D6

protein.

Formulation Liquid in PBS containing 50% glycerol, 0.5% BSA and

0.02% sodium azide.

**Storage** Store at -20°C. Avoid repeated freeze-thaw cycles.

Protein Name Olfactory receptor 4D6

Gene Name OR4D6

Cellular localizationCell membrane; Multi-pass membrane protein.PurificationThe antibody was affinity-purified from rabbit antiserum by affinity-chromatography using

epitope-specific immunogen.

ClonalityPolyclonalConcentration1 mg/mlObserved band36kDHuman Gene ID219983Human Swiss-Prot NumberQ8NGJ1

Alternative Names OR4D6; Olfactory receptor 4D6; Olfactory receptor

OR11-250

Background olfactory receptor family 4 subfamily D member

6(OR4D6) Homo sapiens Olfactory receptors interact with odorant molecules in the nose, to initiate a neuronal response that triggers the perception of a smell. The olfactory receptor




+86-27-59760950 ELKbio@ELKbiotech.com




proteins are members of a large family of G-protein-coupled receptors (GPCR) arising from single coding-exon genes. Olfactory receptors share a 7-transmembrane domain structure with many neurotransmitter and hormone receptors and are responsible for the recognition and G protein-mediated transduction of odorant signals. The olfactory receptor gene family is the largest in the genome. The nomenclature assigned to the olfactory receptor genes and proteins for this organism is independent of other organisms. [provided by RefSeq, Jul 2008],

Western Blot analysis of LOVO cells using Olfactory receptor 4D6 Polyclonal Antibody




Western blot analysis of lysates from LOVO and NIH/3T3 cells, using OR4D6 Antibody. The lane on the right is blocked with the synthesized peptide.



+86-27-59760950







Western blot analysis of the lysates from HT-29 cells using  $\mbox{OR4D6}$  antibody.

