

TFII-I (phospho Tyr248) rabbit pAb

Cat No.: ES5689

For research use only

Overview

Product Name TFII-I (phospho Tyr248) rabbit pAb

Host species Rabbit
Applications WB;ELISA

Species Cross-Reactivity Human; Mouse; Rat

Recommended dilutions Western Blot: 1/500 - 1/2000. ELISA: 1/5000. Not

yet tested in other applications.

Immunogen The antiserum was produced against synthesized

peptide derived from human TFII-I around the phosphorylation site of Tyr248. AA range:214-263

Specificity Phospho-TFII-I (Y248) Polyclonal Antibody detects

endogenous levels of TFII-I protein only when

phosphorylated at Y248.

Formulation Liquid in PBS containing 50% glycerol, 0.5% BSA and

0.02% sodium azide.

Store at -20°C. Avoid repeated freeze-thaw cycles.

Protein Name General transcription factor II-I

Gene Name GTF2I

Cellular localization Cytoplasm . Nucleus . Colocalizes with BTK in the

cytoplasm.

Purification The antibody was affinity-purified from rabbit

antiserum by affinity-chromatography using

epitope-specific immunogen.

Clonality Polyclonal
Concentration 1 mg/ml
Observed band 115kD
Human Gene ID 2969
Human Swiss-Prot Number P78347

Alternative Names GTF2I; BAP135; WBSCR6; General transcription

factor II-I; GTFII-I; TFII-I; Bruton tyrosine kinase-associated protein 135; BAP-135;

BTK-associated protein 135; SRF-Phox1-interacting

protein; SPIN; Williams-Beuren syndrome

chromosomal region




+86-27-59760950 ELKbio@ELKbiotech.com www.elkbiotech.com




Background

general transcription factor IIi(GTF2I) Homo sapiens This gene encodes a phosphoprotein containing six characteristic repeat motifs. The encoded protein binds to the initiator element (Inr) and E-box element in promoters and functions as a regulator of transcription. This locus, along with several other neighboring genes, is deleted in Williams-Beuren syndrome. There are many closely related genes and pseudogenes for this gene on chromosome 7. This gene also has pseudogenes on chromosomes 9, 13, and 21. Alternatively spliced transcript variants encoding multiple isoforms have been observed. [provided by RefSeq, Jul 2013],



Enzyme-Linked Immunosorbent Assay (Phospho-ELISA) for Immunogen Phosphopeptide (Phospho-left) and Non-Phosphopeptide (Phospho-right), using TFII-I (Phospho-Tyr248) Antibody



Western blot analysis of lysates from LOVO cells, using TFII-I (Phospho-Tyr248) Antibody. The lane on the right is blocked with the phospho peptide.

