

hnRNP K (phospho Ser284) rabbit pAb

Cat No.:ES5757

For research use only

Overview

Product Name hnRNP K (phospho Ser284) rabbit pAb

Host species Rabbit

Applications WB;IHC;IF;ELISA

Species Cross-Reactivity Human; Mouse; Rat; Monkey **Recommended dilutions** Western Blot: 1/500 - 1/2000.

Immunohistochemistry: 1/100 - 1/300. ELISA: 1/20000. Not yet tested in other applications.

Immunogen The antiserum was produced against synthesized

peptide derived from human hnRNP K around the phosphorylation site of Ser284. AA range:250-299

Specificity Phospho-hnRNP K (S284) Polyclonal Antibody

detects endogenous levels of hnRNP K protein only

when phosphorylated at S284.

Formulation Liquid in PBS containing 50% glycerol, 0.5% BSA and

0.02% sodium azide.

Storage Store at -20°C. Avoid repeated freeze-thaw cycles.

Protein Name Heterogeneous nuclear ribonucleoprotein K

Gene Name HNRNPK

Cellular localization Cytoplasm . Nucleus, nucleoplasm . Cell projection,

podosome . Recruited to p53/TP53-responsive promoters, in the presence of functional p53/TP53 (PubMed:16360036). In case of ASFV infection, there is a shift in the localization which becomes

predominantl

Purification The antibody was affinity-purified from rabbit

antiserum by affinity-chromatography using

epitope-specific immunogen.

Clonality Polyclonal
Concentration 1 mg/ml
Observed band 51kD
Human Gene ID 3190
Human Swiss-Prot Number P61978

Alternative Names HNRNPK; HNRPK; Heterogeneous nuclear



+86-27-59760950 ELKbio@ELKbiotech.com www.elkbiotech.com




Background

ribonucleoprotein K; hnRNP K; Transformation up-regulated nuclear protein; TUNP This gene belongs to the subfamily of ubiquitously expressed heterogeneous nuclear ribonucleoproteins (hnRNPs). The hnRNPs are RNA binding proteins and they complex with heterogeneous nuclear RNA (hnRNA). These proteins are associated with pre-mRNAs in the nucleus and appear to influence pre-mRNA processing and other aspects of mRNA metabolism and transport. While all of the hnRNPs are present in the nucleus, some seem to shuttle between the nucleus and the cytoplasm. The hnRNP proteins have distinct nucleic acid binding properties. The protein encoded by this gene is located in the nucleoplasm and has three repeats of KH domains that binds to RNAs. It is distinct among other hnRNP proteins in its binding preference; it binds tenaciously to poly(C). This protein is also thought to have a role during cell cycle progession. Several alternatively spliced transcript variants have


3T3 138---100---70---55---40---25---P-hnRNP K (S284) Western Blot analysis of 3T3 cells using Phospho-hnRNP K (S284) Polyclonal Antibody







Immunohistochemical analysis of paraffin-embedded Human brain. Antibody was diluted at 1:100(4° overnight). High-pressure and temperature Tris-EDTA,pH8.0 was used for antigen retrieval. Negetive contrl (right) obtaned from antibody was pre-absorbed by i



Western blot analysis of lysates from NIH/3T3 cells treated with EGF 200ng/ml 30', using hnRNP K (Phospho-Ser284) Antibody. The lane on the right is blocked with the phospho peptide.

