

Trk A (phospho Tyr496) rabbit pAb

Cat No.: ES6425

For research use only

Overview

Product Name Trk A (phospho Tyr496) rabbit pAb

Host species Rabbit

Applications WB;IF;ELISA

Species Cross-Reactivity Human; Mouse; Rat

Recommended dilutions Immunofluorescence: 1/200 - 1/1000. ELISA:

1/5000. Not yet tested in other applications.

Immunogen The antiserum was produced against synthesized

peptide derived from human Trk A around the phosphorylation site of Tyr496. AA range:471-520

Specificity Phospho-Trk A (Y496) Polyclonal Antibody detects

endogenous levels of Trk A protein only when

phosphorylated at Y496.

Formulation Liquid in PBS containing 50% glycerol, 0.5% BSA and

0.02% sodium azide.

Store at -20°C. Avoid repeated freeze-thaw cycles.

Protein Name High affinity nerve growth factor receptor

Gene Name NTRK1

Cellular localization Cell membrane ; Single-pass type I membrane

protein . Early endosome membrane ; Single-pass

type I membrane protein . Late endosome

membrane; Single-pass type I membrane protein.
Recycling endosome membrane; Single-pass type I

membrane protein . Rapidl

Purification The antibody was affinity-purified from rabbit

antiserum by affinity-chromatography using

epitope-specific immunogen.

Clonality Polyclonal
Concentration 1 mg/ml
Observed band 140-180kD
Human Gene ID 4914

Human Gene ID 4914
Human Swiss-Prot Number P04629

Alternative Names NTRK1; MTC; TRK; TRKA; High affinity nerve growth

factor receptor; Neurotrophic tyrosine kinase




+86-27-59760950 ELKbio@ELKbiotech.com

www.elkbiotech.com




Background

receptor type 1; TRK1-transforming tyrosine kinase protein; Tropomyosin-related kinase A; Tyrosine kinase receptor; Tyrosine kinase receptor A; This gene encodes a member of the neurotrophic tyrosine kinase receptor (NTKR) family. This kinase is a membrane-bound receptor that, upon neurotrophin binding, phosphorylates itself and members of the MAPK pathway. The presence of this kinase leads to cell differentiation and may play a role in specifying sensory neuron subtypes. Mutations in this gene have been associated with congenital insensitivity to pain, anhidrosis, self-mutilating behavior, mental retardation and cancer. Alternate transcriptional splice variants of this gene have been found, but only three have been characterized to date. [provided by RefSeq, Jul 2008],



Enzyme-Linked Immunosorbent Assay (Phospho-ELISA) for Immunogen Phosphopeptide (Phospho-left) and Non-Phosphopeptide (Phospho-right), using Trk A (Phospho-Tyr496) Antibody



Immunofluorescence analysis of NIH/3T3 cells, using Trk A (Phospho-Tyr496) Antibody. The picture on the right is blocked with the phospho peptide.

