

Parkin (phospho Ser131) rabbit pAb

Cat No.:ES6478

For research use only

Overview

Product Name	Parkin (phospho Ser131) rabbit pAb	
Host species	Rabbit	
Applications	WB;IHC;IF;ELISA	
Species Cross-Reactivity	Human;Rat;Mouse;	
Recommended dilutions	Immunohistochemistry: 1/100 - 1/300. ELISA:	
	1/20000. Not yet tested in other applications.	
Immunogen	The antiserum was produced against synthesized	
0	peptide derived from human Parkin around the	
	phosphorylation site of Ser131. AA range:101-150	
Specificity	Phospho-Parkin (S131) Polyclonal Antibody detects	
	endogenous levels of Parkin protein only when	
	phosphorylated at \$131.	
Formulation	Liquid in PBS containing 50% glycerol, 0.5% BSA and	
	0.02% sodium azide.	
Storage	Store at -20°C. Avoid repeated freeze-thaw cycles.	
Protein Name	E3 ubiquitin-protein ligase parkin	
Gene Name	PARK2	
Cellular localization	Cytoplasm, cytosol . Nucleus . Endoplasmic	
	reticulum . Mitochondrion . Mitochondrion outer	
	membrane . Cell projection, neuron projection . Cell	
	junction, synapse, postsynaptic density. Cell	
	junction, synapse, presynapse . Mainly localizes in	
	the cytosol	
Purification	The antibody was affinity-purified from rabbit	
	antiserum by affinity-chromatography using	
	epitope-specific immunogen.	
Clonality	Polyclonal	
Concentration	1 mg/ml	
Observed band	51kD	
Human Gene ID	5071	
Human Swiss-Prot Number	O60260	
Alternative Names	PARK2; PRKN; E3 ubiquitin-protein ligase parkin;	
	Parkinson juvenile disease protein 2; Parkinson	



+86-27-59760950

ELKbio@ELKbiotech.com

www.elkbiotech.com

23-2, No.388 Gaoxin 2nd Road, Wuhan East Lake Hi-tech Development Zone, Hubei , P.R.C




Background

disease protein 2

The precise function of this gene is unknown; however, the encoded protein is a component of a multiprotein E3 ubiquitin ligase complex that mediates the targeting of substrate proteins for proteasomal degradation. Mutations in this gene are known to cause Parkinson disease and autosomal recessive juvenile Parkinson disease. Alternative splicing of this gene produces multiple transcript variants encoding distinct isoforms. Additional splice variants of this gene have been described but currently lack transcript support. [provided by RefSeq, Jul 2008],

Western blot analysis of 293T AD293 Hela using p-Parkin (S131) antibody. Antibody was diluted at 1:500



0.600 0.500 0.400 0.300 0.200 0.100 0.000 phosphopeptide non-phosphopeptide I OD 450nm Reading

Enzyme-Linked Immunosorbent Assay (Phospho-ELISA) for Immunogen Phosphopeptide (Phospho-left) and Non-Phosphopeptide (Phospho-right), using Parkin (Phospho-Ser131) Antibody




+86-27-59760950

ELKbio@ELKbiotech.com

www.elkbiotech.com

23-2, No.388 Gaoxin 2nd Road, Wuhan East Lake Hi-tech Development Zone, Hubei , P.R.C





Immunohistochemistry analysis of paraffin-embedded human skeletal muscle, using Parkin (Phospho-Ser131) Antibody. The picture on the right is blocked with the phospho peptide.



+86-27-59760950

ELKbio@ELKbiotech.com

www.elkbiotech.com

23-2, No.388 Gaoxin 2nd Road, Wuhan East Lake Hi-tech Development Zone, Hubei , P.R.C