

## cGKII (phospho Ser126) rabbit pAb

Cat No.: ES6812

For research use only

## Overview

Product Name cGKII (phospho Ser126) rabbit pAb

Host species Rabbit
Applications IHC;IF;ELISA

Species Cross-Reactivity Human; Mouse; Rat

**Recommended dilutions** Immunohistochemistry: 1/100 - 1/300. ELISA:

1/5000. Not yet tested in other applications.

Immunogen The antiserum was produced against synthesized

peptide derived from human PKG2 around the phosphorylation site of Ser126. AA range:92-141

**Specificity** Phospho-cGKII (S126) Polyclonal Antibody detects

endogenous levels of cGKII protein only when

phosphorylated at S126.

Formulation Liquid in PBS containing 50% glycerol, 0.5% BSA and

0.02% sodium azide.

**Store at -20°C.** Avoid repeated freeze-thaw cycles.

**Protein Name** cGMP-dependent protein kinase 2

Gene Name PRKG2

**Cellular localization** Apical cell membrane; Lipid-anchor.

**Purification** The antibody was affinity-purified from rabbit

antiserum by affinity-chromatography using

epitope-specific immunogen.

Clonality Polyclonal Concentration 1 mg/ml

**Observed band** 

Human Gene ID 5593 Human Swiss-Prot Number Q13237

Alternative Names PRKG2; PRKGR2; cGMP-dependent protein kinase 2;

cGK 2; cGK2; cGMP-dependent protein kinase II;

cGKII

**Background** This gene encodes a protein that belongs to the

serine/threonine protein kinase family of proteins.
The encoded protein plays a role in the regulation of fluid balance in the intestine. A similar protein in

+86-27-59760950


ELKbio@ELKbiotech.com

www.elkbiotech.com






mouse is thought to regulate differentiation and proliferation of cells in the colon. Alternate splicing results in multiple transcript variants. [provided by RefSeq, Sep 2013],



Immunohistochemical analysis of paraffin-embedded human tonsil. 1, Antibody was diluted at 1:200(4° overnight). 2, Tris-EDTA,pH9.0 was used for antigen retrieval. 3,Secondary antibody was diluted at 1:200(room temperature, 45min).



Western Blot analysis of various, using primary antibody at 1:1000 dilution. Secondary antibody(catalog#:RS23920) was diluted at 1:10000



+86-27-59760950