



LPAAT- γ rabbit pAb

Cat No.:ES6861

For research use only


Overview

Product Name	LPAAT- γ rabbit pAb
Host species	Rabbit
Applications	WB;ELISA;IHC
Species Cross-Reactivity	Human;Mouse;Rat
Recommended dilutions	WB 1:500-2000;IHC-p 1:50-300; ELISA 2000-20000
Immunogen	The antiserum was produced against synthesized peptide derived from human AGPAT3. AA range:121-170
Specificity	LPAAT- γ Polyclonal Antibody detects endogenous levels of LPAAT- γ protein.
Formulation	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.
Storage	Store at -20°C. Avoid repeated freeze-thaw cycles.
Protein Name	1-acyl-sn-glycerol-3-phosphate acyltransferase gamma
Gene Name	AGPAT3
Cellular localization	Endoplasmic reticulum membrane ; Multi-pass membrane protein . Nucleus envelope .
Purification	The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen.
Clonality	Polyclonal
Concentration	1 mg/ml
Observed band	40kD
Human Gene ID	56894
Human Swiss-Prot Number	Q9NRZ7
Alternative Names	AGPAT3; 1-acyl-sn-glycerol-3-phosphate acyltransferase gamma; 1-acylglycerol-3-phosphate O-acyltransferase 3; 1-AGP acyltransferase 3; 1-AGPAT 3; Lysophosphatidic acid acyltransferase gamma; LPAAT-gamma
Background	The protein encoded by this gene is an acyltransferase that converts lysophosphatidic acid







into phosphatidic acid, which is the second step in the de novo phospholipid biosynthetic pathway. The encoded protein may be an integral membrane protein. Two transcript variants encoding the same protein have been found for this gene. [provided by RefSeq, Jul 2008],



Western blot analysis of lysates from K562 cells, using AGPAT3 Antibody. The lane on the right is blocked with the synthesized peptide.



Western blot analysis of the lysates from COLO205 cells using AGPAT3 antibody.



Immunohistochemical analysis of paraffin-embedded human Gastric adenocarcinoma. 1, Antibody was diluted at 1:200(4° overnight). 2, Tris-EDTA,pH9.0 was used for antigen retrieval. 3,Secondary antibody was diluted at 1:200(room temperature, 45min).

