

AS250 rabbit pAb

Cat No.:ES6875

For research use only

Overview

Product Name AS250 rabbit pAb

Host species Rabbit

Applications WB;IHC;IF;ELISA **Species Cross-Reactivity** Human;Mouse

Recommended dilutions Western Blot: 1/500 - 1/2000.

Immunohistochemistry: 1/100 - 1/300. Immunofluorescence: 1/200 - 1/1000. ELISA:

1/10000. Not yet tested in other applications.

The antiserum was produced against synthesized

Immunogen The antiserum was produced against synthesized

peptide derived from human AS250. AA

range:641-690

Specificity AS250 Polyclonal Antibody detects endogenous

levels of AS250 protein.

Formulation Liquid in PBS containing 50% glycerol, 0.5% BSA and

0.02% sodium azide.

Storage Store at -20°C. Avoid repeated freeze-thaw cycles.

Protein Name Ral GTPase-activating protein subunit alpha-2

Gene Name RALGAPA2 **Cellular localization** Cytoplasm .

Purification The antibody was affinity-purified from rabbit

antiserum by affinity-chromatography using

epitope-specific immunogen.

Clonality Polyclonal
Concentration 1 mg/ml
Observed band 210kD
Human Gene ID 57186
Human Swiss-Prot Number Q2PPJ7

Alternative Names RALGAPA2; C20orf74; KIAA1272; Ral

GTPase-activating protein subunit alpha-2; 250 kDa

substrate of Akt; AS250; p220

Background RALGAPA2 (Ral GTPase Activating Protein Catalytic

Alpha Subunit 2) is a Protein Coding gene. Among its

related pathways are Vesicle-mediated




+86-27-59760950 ELKbio@ELKbiotech.com www.elkbiotech.co




transport and Translocation of GLUT4 to the plasma membrane. GO annotations related to this gene include protein heterodimerization activity and GTPase activator activity. An important paralog of this gene is RAP1GAP.

Immunofluorescence analysis of HeLa cells, using AS250 Antibody. The picture on the right is blocked with the synthesized peptide.



Immunohistochemistry analysis of paraffin-embedded human colon carcinoma tissue, using AS250 Antibody. The picture on the right is blocked with the synthesized peptide.



Western blot analysis of lysates from Jurkat cells, using AS250 Antibody. The lane on the right is blocked with the synthesized peptide.

