

ARHGAP9 rabbit pAb

Cat No.:ES7164

For research use only

Overview

Product Name ARHGAP9 rabbit pAb

Host species Rabbit

Applications WB;IHC;IF;ELISA Species Cross-Reactivity Human;Rat;Mouse;

Recommended dilutions Western Blot: 1/500 - 1/2000.

Immunohistochemistry: 1/100 - 1/300. ELISA: 1/40000. Not yet tested in other applications.

Immunogen The antiserum was produced against synthesized

peptide derived from human RHG9. AA

range:220-269

Specificity ARHGAP9 Polyclonal Antibody detects endogenous

levels of ARHGAP9 protein.

Formulation Liquid in PBS containing 50% glycerol, 0.5% BSA and

0.02% sodium azide.

Storage Store at -20°C. Avoid repeated freeze-thaw cycles.

Protein Name Rho GTPase-activating protein 9

Gene Name ARHGAP9
Cellular localization cytosol,

Purification The antibody was affinity-purified from rabbit

antiserum by affinity-chromatography using

epitope-specific immunogen.

ClonalityPolyclonalConcentration1 mg/mlObserved band90kDHuman Gene ID64333Human Swiss-Prot NumberQ9BRR9

Alternative Names ARHGAP9; Rho GTPase-activating protein 9;

Rho-type GTPase-activating protein 9

Background This gene encodes a member of the Rho-GAP family

of GTPase activating proteins. The protein has substantial GAP activity towards several Rho-family GTPases in vitro, converting them to an inactive GDP-bound state. It is implicated in regulating




+86-27-59760950 ELKbio@ELKbiotech.com




-- 117 -- 85 -- 48 -- 34 -- 26 -- 19 (kD) adhesion of hematopoietic cells to the extracellular matrix. Multiple transcript variants encoding different isoforms have been found for this gene. [provided by RefSeq, Jul 2008],

Western blot analysis of lysates from LOVO cells, using RHG9 Antibody. The lane on the right is blocked with the synthesized peptide.



Western blot analysis of the lysates from COLO205 cells using RHG9 antibody.



+86-27-59760950

Immunohistochemical analysis of paraffin-embedded human Gastric adenocarcinoma. 1, Antibody was diluted at 1:200(4° overnight). 2, Tris-EDTA,pH9.0 was used for antigen retrieval. 3,Secondary antibody was diluted at 1:200(room temperature, 45min).

