

POLR1E rabbit pAb

Cat No.:ES7174

For research use only

Overview

Product Name	POLR1E rabbit pAb
Host species	Rabbit
Applications	WB;IHC;IF;ELISA
Species Cross-Reactivity	Human;Rat;Mouse;
Recommended dilutions	Western Blot: 1/500 - 1/2000.
	Immunohistochemistry: 1/100 - 1/300. ELISA:
	1/10000. Not yet tested in other applications.
Immunogen	Synthesized peptide derived from the Internal
	region of human POLR1E.
Specificity	POLR1E Polyclonal Antibody detects endogenous
	levels of POLR1E protein.
Formulation	Liquid in PBS containing 50% glycerol, 0.5% BSA and
	0.02% sodium azide.
Storage	Store at -20°C. Avoid repeated freeze-thaw cycles.
Protein Name	DNA-directed RNA polymerase I subunit RPA49
Gene Name	POLR1E
Cellular localization	Nucleus, nucleolus .
Purification	The antibody was affinity-purified from rabbit
	antiserum by affinity-chromatography using
	epitope-specific immunogen.
Clonality	Polyclonal
Concentration	1 mg/ml
Observed band	49kD
Human Gene ID	64425
Human Swiss-Prot Number	Q9GZS1
Alternative Names	POLR1E; PAF53; PRAF1; DNA-directed RNA
	polymerase I subunit RPA49; RNA polymerase I
	subunit A49; DNA-directed RNA polymerase I
	subunit E; RNA polymerase I-associated factor 1;
	RNA polymerase I-associated factor 53
Background	function:DNA-dependent RNA polymerase catalyzes
	the transcription of DNA into RNA using the four
	ribonucleoside triphosphates as substrates.



+86-27-59760950

ELKbio@ELKbiotech.com

www.elkbiotech.com

23-2, No.388 Gaoxin 2nd Road, Wuhan East Lake Hi-tech Development Zone, Hubei , P.R.C



Component of RNA polymerase I which synthesizes ribosomal RNA precursors. Appears to be involved in the formation of the initiation complex at the promoter by mediating the interaction between Pol I and UBTF/UBF.,similarity:Belongs to the eukaryotic RPA49/POLR1E RNA polymerase subunit family.,subunit:Component of the RNA polymerase I (Pol I) complex consisting of at least 13 subunits (By similarity). Interacts with PAF49/CD3EAP. Also binds UBTF/UBF.,

Immunohistochemical analysis of paraffin-embedded human brain. 1, Tris-EDTA,pH9.0 was used for antigen retrieval. 2 Antibody was diluted at 1:200(4° overnight.3,Secondary antibody was diluted at 1:200(room temperature, 45min).





+86-27-59760950

ELKbio@ELKbiotech.com

www.elkbiotech.com

23-2, No.388 Gaoxin 2nd Road, Wuhan East Lake Hi-tech Development Zone, Hubei , P.R.C