

# Endophilin I rabbit pAb

Cat No.:ES7184

For research use only


## Overview

|                          |  |
|--------------------------|--|
| Product Name             | Endophilin I rabbit pAb  |
| Host species             | Rabbit   |
| Applications             | WB;ELISA   |
| Species Cross-Reactivity | Human;Mouse;Rat  |
| Recommended dilutions    | Western Blot: 1/500 - 1/2000. ELISA: 1/10000. Not yet tested in other applications.  |
| Immunogen                | Synthesized peptide derived from Endophilin I . at AA range: 30-110  |
| Specificity              | Endophilin I Polyclonal Antibody detects endogenous levels of Endophilin I protein.  |
| Formulation              | Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.  |
| Storage                  | Store at -20°C. Avoid repeated freeze-thaw cycles.   |
| Protein Name             | Endophilin-A1  |
| Gene Name                | SH3GL2   |
| Cellular localization    | Cytoplasm . Membrane ; Peripheral membrane protein . Early endosome . Cell junction, synapse, presynapse .   |
| Purification             | The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen.  |
| Clonality                | Polyclonal   |
| Concentration            | 1 mg/ml  |
| Observed band            | 39kD   |
| Human Gene ID            | 6456   |
| Human Swiss-Prot Number  | Q99962   |
| Alternative Names        | SH3GL2; CNSA2; SH3D2A; Endophilin-A1; EEN-B1; Endophilin-1; SH3 domain protein 2A; SH3 domain-containing GRB2-like protein 2   |
| Background               | domain:An N-terminal amphipathic helix, the BAR domain and a second amphipathic helix inserted into helix 1 of the BAR domain (N-BAR domain) induce membrane curvature and bind curved |





membranes. The BAR domain dimer forms a rigid crescent shaped bundle of helices with the pair of second amphipathic helices protruding towards the membrane-binding surface.,function:Implicated in synaptic vesicle endocytosis. May recruit other proteins to membranes with high curvature.,miscellaneous:HeLa cells expressing the N-BAR domain of SH3GL2 show tubulation of the plasma membrane. The N-BAR domain binds liposomes and induces formation of tubules from liposomes. The N-terminal amphipathic helix is required for liposome binding. The second amphipathic helix enhances liposome tubulation.,similarity:Belongs to the endophilin family.,similarity:Contains 1 BAR domain.,similarity:Contains 1 SH3 domain.,subcellular location:Concentrated in presynaptic nerve terminals in neurons.,subunit:Monomer; in cytoplasm. Homodimer; when associated with membranes (By similarity). Interacts with SYNJ1 and DNM1. Interacts with MAP4K3; the interaction appears to regulate MAP4K3-mediated JNK activation. Interacts with PDCD6IP.,tissue specificity:Brain, mostly in frontal cortex. Expressed at high level in fetal cerebellum.,



Western blot analysis of SH3GL2 Antibody. The lane on the right is blocked with the SH3GL2 peptide.

