

EMR3 rabbit pAb

Cat No.: ES7772

For research use only

Overview

Product Name EMR3 rabbit pAb

Host species Rabbit

Applications WB;IF;ELISA

Species Cross-Reactivity Human; Rat; Mouse;

Recommended dilutions Western Blot: 1/500 - 1/2000. Immunofluorescence:

1/200 - 1/1000. ELISA: 1/10000. Not yet tested in

other applications.

Immunogen The antiserum was produced against synthesized

peptide derived from human EMR3. AA

range:603-652

Specificity EMR3 Polyclonal Antibody detects endogenous

levels of EMR3 protein.

Formulation Liquid in PBS containing 50% glycerol, 0.5% BSA and

0.02% sodium azide.

Storage Store at -20°C. Avoid repeated freeze-thaw cycles.

Protein Name EGF-like module-containing mucin-like hormone

receptor-like 3

Gene Name EMR3

Cell ular localization Cell membrane; Multi-pass membrane protein .;

[Isoform 3]: Secreted.

Purification The antibody was affinity-purified from rabbit

antiserum by affinity-chromatography using

epitope-specific immunogen.

Clonality Polyclonal
Concentration 1 mg/ml
Observed band 70kD
Human Gene ID 84658
Human Swiss-Prot Number Q9BY15

Alternative Names EMR3; EGF-like module-containing mucin-like

hormone receptor-like 3; EGF-like module receptor

3

Background This gene encodes a member of the class B


seven-span transmembrane (TM7) receptor family



+86-27-59760950 ELKbio@ELKbiotech.com www.elkbiotech.com



expressed predominantly by cells of the immune system. Family members are characterized by an extended extracellular region with a variable number of N-terminal epidermal growth factor (EGF)-like domains coupled to a TM7 domain via a mucin-like spacer domain. This gene is closely linked to the gene encoding egf-like molecule containing mucin-like hormone receptor 2 on chromosome 19. This protein may play a role in myeloid-myeloid interactions during immune and inflammatory responses. Alternative splicing results in multiple transcript variants encoding different isoforms. [provided by RefSeq, Jan 2014],



Western blot analysis of lysates from Jurkat cells, using EMR3 Antibody. The lane on the right is blocked with the synthesized peptide.

