

Claudin-6 (phospho Tyr219) rabbit pAb

Cat No.:ES7950

For research use only

Overview

Product Name	Claudin-6 (phospho Tyr219) rabbit pAb
Host species	Rabbit
Applications	WB;IHC;IF;ELISA
Species Cross-Reactivity	Human; Mouse
Recommended dilutions	Western Blot: 1/500 - 1/2000.
	Immunohistochemistry: 1/100 - 1/300. ELISA:
	1/5000. Not yet tested in other applications.
Immunogen	The antiserum was produced against synthesized
	peptide derived from human Claudin 6 around the
	phosphorylation site of Tyr219. AA range:171-220
Specificity	Phospho-Claudin-6 (Y219) Polyclonal Antibody
	detects endogenous levels of Claudin-6 protein only
	when phosphorylated at Y219.
Formulation	Liquid in PBS containing 50% glycerol, 0.5% BSA and
	0.02% sodium azide.
Storage	Store at -20°C. Avoid repeated freeze-thaw cycles.
Protein Name	Claudin-6
Gene Name	CLDN6
Cellular localization	Cell junction, tight junction . Cell membrane ;
	Multi-pass membrane protein .
Purification	The antibody was affinity-purified from rabbit
	antiserum by affinity-chromatography using
	epitope-specific immunogen.
Clonality	Polyclonal
Concentration	1 mg/ml
Observed band	23kD
Human Gene ID	9074
Human Swiss-Prot Number	P56747
Alternative Names	CLDN6; Claudin-6; Skullin
Background	Tight junctions represent one mode of cell-to-cell
	adhesion in epithelial or endothelial cell sheets,
	forming continuous seals around cells and serving as
	a physical barrier to prevent solutes and water from



+86-27-59760950


ELKbio@ELKbiotech.com

www.elkbiotech.com

23-2, No.388 Gaoxin 2nd Road, Wuhan East Lake Hi-tech Development Zone, Hubei , P.R.C



passing freely through the paracellular space. These junctions are comprised of sets of continuous networking strands in the outwardly facing cytoplasmic leaflet, with complementary grooves in the inwardly facing extracytoplasmic leaflet. This gene encodes a component of tight junction strands, which is a member of the claudin family. The protein is an integral membrane protein and is one of the entry cofactors for hepatitis C virus. The gene methylation may be involved in esophageal tumorigenesis. This gene is adjacent to another family member CLDN9 on chromosome 16.[provided by RefSeq, Aug 2010],



Western blot analysis of lysates from RAW264.7 cells treated with UV 5', using Claudin 6 (Phospho-Tyr219) Antibody. The lane on the right is blocked with the phospho peptide.

retrieval. 3,Seco temperature, 4

Immunohistochemical analysis of paraffin-embedded human tonsil. 1, Antibody was diluted at 1:200(4° overnight). 2, Tris-EDTA,pH9.0 was used for antigen retrieval. 3,Secondary antibody was diluted at 1:200(room temperature, 45min).



+86-27-59760950

ELKbio@ELKbiotech.com

www.elkbiotech.com

23-2, No.388 Gaoxin 2nd Road, Wuhan East Lake Hi-tech Development Zone, Hubei , P.R.C