

## Mox1 rabbit pAb

Cat No.: ES8635

For research use only

## Overview

Product Name Mox1 rabbit pAb

Host species Rabbit

Applications WB;IHC;IF;ELISA Species Cross-Reactivity Human;Mouse;Rat

Recommended dilutions WB 1:500-2000,IHC-p 1:500-200, ELISA

1:10000-20000

**Immunogen** Synthetic peptide from human protein at AA range:

210-260

**Specificity** The antibody detects endogenous Mox1

Formulation Liquid in PBS containing 50% glycerol, 0.5% BSA and

0.02% sodium azide.

Storage Store at -20°C. Avoid repeated freeze-thaw cycles.

Protein Name NADPH oxidase 1 (NOX-1) (EC 1.-.-.) (Mitogenic oxidase 1) (MOX-1) (NADPH mitogenic

oxidase 1) (MOX-1) (NADH/NADPH mitogenic

oxidase subunit P65-MOX) (NOH-1)

Gene Name NOX1 MOX1 NOH1

**Cellular localization** Cell projection, invadopodium membrane;

Multi-pass membrane protein . Cell membrane .

**Purification** The antibody was affinity-purified from rabbit

antiserum by affinity-chromatography using

epitope-specific immunogen.

Clonality Polyclonal
Concentration 1 mg/ml
Observed band 65kD
Human Gene ID 27035
Human Swiss-Prot Number Q9Y5S8

Alternative Names NADPH oxidase 1 (NOX-1;EC 1.-.-.;Mitogenic

oxidase 1;MOX-1;NADH/NADPH mitogenic oxidase

subunit P65-MOX;NOH-1)

**Background** This gene encodes a member of the NADPH oxidase


family of enzymes responsible for the catalytic one-electron transfer of oxygen to generate superoxide or hydrogen peroxide. Alternatively




+86-27-59760950 ELKbio@ELKbiotech.com www.elkbiote




spliced transcript variants encoding multiple isoforms have been observed for this gene. [provided by RefSeq, Nov 2012],



Western blot analysis of 293T Hela VEC KB mouse-kidney 3T3 lysate, antibody was diluted at 2000. Secondary antibody(catalog#:RS0002) was diluted at 1:20000



Immunohistochemical analysis of paraffin-embedded human-liver, antibody was diluted at 1:200



Immunohistochemical analysis of paraffin-embedded human-liver, antibody was diluted at 1:200





Immunohistochemical analysis of paraffin-embedded human-stomach, antibody was diluted at 1:200



+86-27-59760950