

Tubulin α rabbit pAb

Cat No.: ES8766

For research use only

Overview

Product Name Tubulin α rabbit pAb

Host species Rabbit WB;ELISA **Applications**

Species Cross-Reactivity Human;Rat;Mouse

Recommended dilutions WB 1:500-2000, ELISA 1:10000-20000

Immunogen Synthetic peptide from human protein at AA range:

382-451

The antibody detects endogenous Tubulin α Specificity

Formulation Liquid in PBS containing 50% glycerol, 0.5% BSA and

0.02% sodium azide.

Storage Store at -20°C. Avoid repeated freeze-thaw cycles.

Protein Name Tubulin α TUBA1A/B Gene Name

Cellular localization Cytoplasm, cytoskeleton.

Purification The antibody was affinity-purified from rabbit

antiserum by affinity-chromatography using

epitope-specific immunogen.

Clonality Polyclonal Concentration 1 mg/ml Observed band 55kD **Human Gene ID** 7846

Human Swiss-Prot Number

Alternative Names

Q71U36/P68363


Background Microtubules of the eukaryotic cytoskeleton

perform essential and diverse functions and are composed of a heterodimer of alpha and beta tubulins. The genes encoding these microtubule constituents belong to the tubulin superfamily, which is composed of six distinct families. Genes from the alpha, beta and gamma tubulin families are found in all eukaryotes. The alpha and beta tubulins represent the major components of microtubules, while gamma tubulin plays a critical role in the



+86-27-59760950 ELKbio@ELKbiotech.com





nucleation of microtubule assembly. There are multiple alpha and beta tubulin genes, which are highly conserved among species. This gene encodes alpha tubulin and is highly similar to the mouse and rat Tuba1 genes. Northern blotting studies have shown that the gene expression is predominantly found in morphologically differentiated neurologic cells. This gene is one of three alpha-tubulin genes in a cluster on chromosome 12q.

Western blot analysis of Hela lysate, antibody was diluted at 2000. Secondary antibody(catalog#:RS0002) was diluted at 1:20000

