

GLI-1 rabbit pAb

Cat No.: ES8816

For research use only

Overview

Product Name GLI-1 rabbit pAb

Host species Rabbit
Applications WB;ELISA
Species Cross-Reactivity Human;Mouse

Recommended dilutions WB 1:500-2000, ELISA 1:10000-20000

Immunogen Synthesized peptide derived from human GLI-1. at

AA range: 460-490

Specificity This antibody detects endogenous levels of GLI-1 **Formulation** Liquid in PBS containing 50% glycerol, 0.5% BSA and

0.02% sodium azide.

Storage Store at -20° C. Avoid repeated freeze-thaw cycles.

Protein Name GLI-1 Gene Name GLI1 GLI

Cellular localization Cytoplasm . Nucleus . Tethered in the cytoplasm by

binding to SUFU (PubMed:10806483). Activation and translocation to the nucleus is promoted by interaction with STK36 (PubMed:10806483). Phosphorylation by ULK3 may promote nuclear localization (PubMed:19878745). Translocation to the nucleus is promoted by interaction with ZIC1 (PubMed:11238441)..; [Isoform 2]: Cytoplasm.

Nucleus.

Purification The antibody was affinity-purified from rabbit

antiserum by affinity-chromatography using

epitope-specific immunogen.

ClonalityPolyclonalConcentration1 mg/mlObserved band120kDHuman Gene ID2735Human Swiss-Prot NumberP08151

Alternative Names Zinc finger protein GLI1 (Glioma-associated

oncogene) (Oncogene GLI)

Background This gene encodes a member of the Kruppel family



+86-27-59760950 ELKbio@ELKbiotech.com www.elkbiotech.com



of zinc finger proteins. The encoded transcription factor is activated by the sonic hedgehog signal transduction cascade and regulates stem cell proliferation. The activity and nuclear localization of this protein is negatively regulated by p53 in an inhibitory loop. Multiple transcript variants encoding different isoforms have been found for this gene. [provided by RefSeq, May 2009],



+86-27-59760950