



AKAP9 rabbit pAb

Cat No.:ES9089

For research use only

Overview

Product Name	AKAP9 rabbit pAb
Host species	Rabbit
Applications	IHC;IF
Species Cross-Reactivity	Human;Rat;Mouse;
Recommended dilutions	IHC-p 1:50-300
Immunogen	Synthesized peptide derived from human protein . at AA range: 2610-2690
Specificity	AKAP9 Polyclonal Antibody detects endogenous levels of protein.
Formulation	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.
Storage	Store at -20°C. Avoid repeated freeze-thaw cycles.
Protein Name	A-kinase anchor protein 9 (AKAP-9) (A-kinase anchor protein 350 kDa) (AKAP 350) (hgAKAP 350) (A-kinase anchor protein 450 kDa) (AKAP 450) (AKAP 120-like protein) (Centrosome- and Golgi-localized PKN-a
Gene Name	AKAP9 AKAP350 AKAP450 KIAA0803
Cellular localization	Golgi apparatus . Cytoplasm . Cytoplasm, cytoskeleton, microtubule organizing center, centrosome . Cytoplasmic in parietal cells (PubMed:9915845). Recruited to the Golgi apparatus by GM130/GOLGA2 (PubMed:25657325). Localization at the centrosome versus Go
Purification	The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen.
Clonality	Polyclonal
Concentration	1 mg/ml
Observed band	430kD
Human Gene ID	10142
Human Swiss-Prot Number	Q99996
Alternative Names	
Background	The A-kinase anchor proteins (AKAPs) are a group of





structurally diverse proteins which have the common function of binding to the regulatory subunit of protein kinase A (PKA) and confining the holoenzyme to discrete locations within the cell. This gene encodes a member of the AKAP family. Alternate splicing of this gene results in at least two isoforms that localize to the centrosome and the Golgi apparatus, and interact with numerous signaling proteins from multiple signal transduction pathways. These signaling proteins include type II protein kinase A, serine/threonine kinase protein kinase N, protein phosphatase 1, protein phosphatase 2a, protein kinase C-epsilon and phosphodiesterase 4D3. [provided by RefSeq, Aug 2008],

Immunohistochemical analysis of paraffin-embedded human tonsil. 1, Antibody was diluted at 1:200(4° overnight). 2, Tris-EDTA,pH9.0 was used for antigen retrieval. 3,Secondary antibody was diluted at 1:200(room temperature, 30min).

