

MAGB1 rabbit pAb

Cat No.: ES9202

For research use only

Overview

Product Name MAGB1 rabbit pAb

Host species Rabbit
Applications WB;ELISA

Species Cross-Reactivity Human; Rat; Mouse;

Recommended dilutions WB 1:500-2000 ELISA 1:5000-20000

Immunogen Synthesized peptide derived from human protein.

at AA range: 70-150

Specificity MAGB1 Polyclonal Antibody detects endogenous

levels of protein.

Formulation Liquid in PBS containing 50% glycerol, 0.5% BSA and

0.02% sodium azide.

Storage Store at -20°C. Avoid repeated freeze-thaw cycles.

Protein Name Melanoma-associated antigen B1 (Cancer/testis

antigen 3.1) (CT3.1) (DSS-AHC critical interval MAGE

superfamily 10) (DAM10) (MAGE-B1 antigen)

(MAGE-XP antigen)

Gene Name MAGEB1 MAGEL1 MAGEXP

Cellular localization

Purification The antibody was affinity-purified from rabbit

antiserum by affinity-chromatography using

epitope-specific immunogen.

Clonality Polyclonal
Concentration 1 mg/ml
Observed band 38kD
Human Gene ID 4112
Human Swiss-Prot Number P43366

Alternative Names

Background This gene is a member of the MAGEB gene family.

The members of this family have their entire coding sequences located in the last exon, and the encoded proteins show 50 to 68% sequence identity to each other. The promoters and first exons of the MAGEB genes show considerable variability, suggesting that




+86-27-59760950 ELKbio@ELKbiotech.com

www.elkbiotech.com



the existence of this gene family enables the same function to be expressed under different transcriptional controls. This gene is localized in the DSS (dosage-sensitive sex reversal) critical region, and expressed in testis and in a significant fraction of tumors of various histological types. This gene and other MAGEB members are clustered on chromosome Xp22-p21. Multiple alternatively spliced transcript variants encoding the same protein have been found for this gene, however, the full length nature of some variants has not been defined. [provided by RefSeq, Jul 2008],

Western blot analysis of lysates from U2OS cells, primary antibody was diluted at 1:1000, 4° over night



+86-27-59760950

