

PER1 rabbit pAb

Cat No.:ES9274

For research use only

Overview

| Product Name | PER1 rabbit pAb | |
|--------------------------|--|--|
| Host species | Rabbit | |
| Applications | WB;ELISA | |
| Species Cross-Reactivity | Human;Mouse;Rat | |
| Recommended dilutions | WB 1:500-2000 ELISA 1:5000-20000 | |
| Immunogen | Synthesized peptide derived from human protein . at | |
| C | AA range: 1110-1190 | |
| Specificity | PER1 Polyclonal Antibody detects endogenous levels | |
| | of protein. | |
| Formulation | Liquid in PBS containing 50% glycerol, 0.5% BSA and | |
| | 0.02% sodium azide. | |
| Storage | Store at -20 $^\circ\!\mathrm{C}$. Avoid repeated freeze-thaw cycles. | |
| Protein Name | Period circadian protein homolog 1 (hPER1) | |
| | (Circadian clock protein PERIOD 1) (Circadian | |
| | pacemaker protein Rigui) | |
| Gene Name | PER1 KIAA0482 PER RIGUI | |
| Cellular localization | Nucleus. Cytoplasm. Nucleocytoplasmic shuttling is | |
| | effected by interaction with other circadian core | |
| | oscillator proteins and/or by phosphorylation. | |
| | Retention of PER1 in the cytoplasm occurs through | |
| | PER1-PER2 heterodimer formation. Translocate to | |
| | the nucleus after phosphorylation by CSNK1D or | |
| | CSNK1E. Also translocated to the nucleus by CRY1 or | |
| | CRY2 (By similarity) | |
| Purification | The antibody was affinity-purified from rabbit | |
| | antiserum by affinity-chromatography using | |
| | epitope-specific immunogen. | |
| Clonality | Polyclonal | |
| Concentration | 1 mg/ml | |
| Observed band | 141kD | |
| Human Gene ID | 5187 | |
| Human Swiss-Prot Number | 015534 | |
| Alternative Names | | |



+86-27-59760950

ELKbio@ELKbiotech.com

www.elkbiotech.com

23-2, No.388 Gaoxin 2nd Road, Wuhan East Lake Hi-tech Development Zone, Hubei , P.R.C



Background

This gene is a member of the Period family of genes and is expressed in a circadian pattern in the suprachiasmatic nucleus, the primary circadian pacemaker in the mammalian brain. Genes in this family encode components of the circadian rhythms of locomotor activity, metabolism, and behavior. This gene is upregulated by CLOCK/ARNTL heterodimers but then represses this upregulation in a feedback loop using PER/CRY heterodimers to interact with CLOCK/ARNTL. Polymorphisms in this gene may increase the risk of getting certain cancers. Alternative splicing has been observed in this gene; however, these variants have not been fully described. [provided by RefSeq, Jan 2014],



+86-27-59760950

ELKbio@ELKbiotech.com

www.elkbiotech.com

23-2, No.388 Gaoxin 2nd Road, Wuhan East Lake Hi-tech Development Zone, Hubei , P.R.C