

## **ADHX** rabbit pAb

Cat No.: ES9372

For research use only

## Overview

Product Name ADHX rabbit pAb

Host species Rabbit
Applications WB;ELISA

Species Cross-Reactivity Human; Mouse; Rat

Recommended dilutions WB 1:500-2000 ELISA 1:5000-20000

**Immunogen** Synthesized peptide derived from human protein .

at AA range: 280-360

**Specificity** ADHX Polyclonal Antibody detects endogenous

levels of protein.

Formulation Liquid in PBS containing 50% glycerol, 0.5% BSA and

0.02% sodium azide.

Storage Store at -20°C. Avoid repeated freeze-thaw cycles.

Protein Name Alcohol dehydrogenase class-3 (EC 1.1.1.1) (Alcohol

dehydrogenase 5) (Alcohol dehydrogenase class chi

chain) (Alcohol dehydrogenase class-III) (Glutathione-dependent formaldehyde

dehydrogenase) (FALDH

Gene Name ADH5 ADHX FDH Cellular localization Cytoplasm .

**Purification** The antibody was affinity-purified from rabbit

antiserum by affinity-chromatography using

epitope-specific immunogen.

Clonality Polyclonal
Concentration 1 mg/ml
Observed band 41kD
Human Gene ID 128
Human Swiss-Prot Number P11766

**Alternative Names** 

**Background** This gene encodes a member of the alcohol

dehydrogenase family. Members of this family metabolize a wide variety of substrates, including

ethanol, retinol, other aliphatic alcohols,

hydroxysteroids, and lipid peroxidation products.




+86-27-59760950 ELKbio@ELKbiotech.com

www.elkbiotech.com



The encoded protein forms a homodimer. It has virtually no activity for ethanol oxidation, but exhibits high activity for oxidation of long-chain primary alcohols and for oxidation of S-hydroxymethyl-glutathione, a spontaneous adduct between formaldehyde and glutathione. This enzyme is an important component of cellular metabolism for the elimination of formaldehyde, a potent irritant and sensitizing agent that causes lacrymation, rhinitis, pharyngitis, and contact dermatitis. The human genome contains several non-transcribed pseudogenes related to this gene. [provided by RefSeq, Oct 2008],

Western blot analysis of lysates from A431 cells, primary antibody was diluted at 1:1000, 4° over night



+86-27-59760950

