



GNA13 rabbit pAb

Cat No.:ES9699

For research use only


Overview

Product Name	GNA13 rabbit pAb
Host species	Rabbit
Applications	WB;ELISA
Species Cross-Reactivity	Human;Mouse
Recommended dilutions	WB 1:500-2000 ELISA 1:5000-20000
Immunogen	Synthesized peptide derived from part region of human protein
Specificity	GNA13 Polyclonal Antibody detects endogenous levels of protein.
Formulation	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.
Storage	Store at -20°C. Avoid repeated freeze-thaw cycles.
Protein Name	Guanine nucleotide-binding protein subunit alpha-13 (G alpha-13) (G-protein subunit alpha-13)
Gene Name	GNA13
Cellular localization	Cell membrane ; Lipid-anchor . Melanosome . Cytoplasm . Nucleus . Identified by mass spectrometry in melanosome fractions from stage I to stage IV (PubMed:17081065). Detected in the cytoplasm of Leydig cells and in the seminiferous epithelium, including d
Purification	The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen.
Clonality	Polyclonal
Concentration	1 mg/ml
Observed band	41kD
Human Gene ID	10672
Human Swiss-Prot Number	Q14344
Alternative Names	
Background	function:Guanine nucleotide-binding proteins (G proteins) are involved as modulators or transducers in various transmembrane signaling





systems.,PTM:Palmitoylation is critical for proper membrane localization and signaling.,PTM:Phosphorylation on Thr-203 by PKA destabilizes the heterotrimer of alpha, beta and gamma, and inhibits Rho activation.,similarity:Belongs to the G-alpha family. G(12) subfamily.,subcellular location:Identified by mass spectrometry in melanosome fractions from stage I to stage IV.,subunit:G proteins are composed of 3 units; alpha, beta and gamma. The alpha chain contains the guanine nucleotide binding site. Interacts with UBXD5.,



Western blot analysis of lysates from K562 cells, primary antibody was diluted at 1:1000, 4° over night

