

MYP0 rabbit pAb

Cat No.: ES9836

For research use only

Overview

Product Name MYP0 rabbit pAb

Host species Rabbit
Applications WB;ELISA

Species Cross-Reactivity Human; Mouse; Rat

Recommended dilutions WB 1:500-2000 ELISA 1:5000-20000

Immunogen Synthesized peptide derived from human protein .

at AA range: 50-130

Specificity MYPO Polyclonal Antibody detects endogenous

levels of protein.

Formulation Liquid in PBS containing 50% glycerol, 0.5% BSA and

0.02% sodium azide.

Storage Store at -20°C. Avoid repeated freeze-thaw cycles.

Protein Name Myelin protein P0 (Myelin peripheral protein) (MPP)

(Myelin protein zero)

Gene Name MPZ

Cellular localization Cell membrane ; Single-pass type I membrane

protein.; [Isoform L-MPZ]: Myelin membrane;

Single-pass type I membrane protein.

Purification The antibody was affinity-purified from rabbit

antiserum by affinity-chromatography using

epitope-specific immunogen.

Clonality Polyclonal
Concentration 1 mg/ml
Observed band 27kD
Human Gene ID 4359
Human Swiss-Prot Number P25189

Alternative Names

Background This gene is specifically expressed in Schwann cells

of the peripheral nervous system and encodes a type I transmembrane glycoprotein that is a major structural protein of the peripheral myelin sheath. The encoded protein contains a large hydrophobic extracellular domain and a smaller basic intracellular




+86-27-59760950 ELKbio@ELKbiotech.com

www.elkbiotech.com



domain, which are essential for the formation and stabilization of the multilamellar structure of the compact myelin. Mutations in this gene are associated with autosomal dominant form of Charcot-Marie-Tooth disease type 1 (CMT1B) and other polyneuropathies, such as Dejerine-Sottas syndrome (DSS) and congenital hypomyelinating neuropathy (CHN). A recent study showed that two isoforms are produced from the same mRNA by use of alternative in-frame translation termination codons via a stop codon readthrough mechanism. [provided by RefSeq, Oct 2015],

Western blot analysis of lysates from DU145 cells, primary antibody was diluted at 1:1000, 4° over night



+86-27-59760950

