

NOG2 rabbit pAb

Cat No.:ES9952

For research use only

Overview

| Product Name | NOG2 rabbit pAb |
|--------------------------|--|
| Host species | Rabbit |
| Applications | WB;ELISA |
| Species Cross-Reactivity | Human;Mouse |
| Recommended dilutions | WB 1:500-2000 ELISA 1:5000-20000 |
| Immunogen | Synthesized peptide derived from human protein . at |
| | AA range: 670-750 |
| Specificity | NOG2 Polyclonal Antibody detects endogenous |
| | levels of protein. |
| Formulation | Liquid in PBS containing 50% glycerol, 0.5% BSA and |
| | 0.02% sodium azide. |
| Storage | Store at -20 $^\circ\!\mathrm{C}$. Avoid repeated freeze-thaw cycles. |
| Protein Name | Nucleolar GTP-binding protein 2 (Autoantigen |
| | NGP-1) |
| Gene Name | GNL2 NGP1 |
| Cellular localization | Nucleus, nucleolus . |
| Purification | The antibody was affinity-purified from rabbit |
| | antiserum by affinity-chromatography using |
| | epitope-specific immunogen. |
| Clonality | Polyclonal |
| Concentration | 1 mg/ml |
| Observed band | 80kD |
| Human Gene ID | 29889 |
| Human Swiss-Prot Number | Q13823 |
| Alternative Names | |
| Background | function:GTPase that associates with pre-60S |
| | ribosomal subunits in the nucleolus and is required |
| | for their nuclear export and |
| | maturation., similarity: Belongs to the MMR1/HSR1 |
| | GTP-binding protein family. NOG2 subfamily., tissue |
| | specificity:Ubiquitously expressed at relatively low |
| | levels in all human tissues tested, with the highest |
| | level of expression in the testes., |



+86-27-59760950

ELKbio@ELKbiotech.com

www.elkbiotech.com

23-2, No.388 Gaoxin 2nd Road, Wuhan East Lake Hi-tech Development Zone, Hubei , P.R.C