

RFIP5 rabbit pAb

Cat No.:ES10079

For research use only

Overview

Product Name	RFIP5 rabbit pAb
Host species	Rabbit
Applications	WB;ELISA
Species Cross-Reactivity	Human;Mouse
Recommended dilutions	WB 1:500-2000 ELISA 1:5000-20000
Immunogen	Synthesized peptide derived from human protein . at AA range: 570-650
Specificity	RFIP5 Polyclonal Antibody detects endogenous levels of protein.
Formulation	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.
Storage	Store at -20°C . Avoid repeated freeze-thaw cycles.
Protein Name	Rab11 family-interacting protein 5 (Rab11-FIP5) (Gamma-SNAP-associated factor 1) (Gaf-1) (Phosphoprotein pp75) (Rab11-interacting protein Rip11)
Gene Name	RAB11FIP5 GAF1 KIAA0857 RIP11
Cellular localization	Cytoplasm. Recycling endosome membrane; Peripheral membrane protein. Early endosome membrane ; Peripheral membrane protein . Golgi apparatus membrane ; Peripheral membrane protein . Cytoplasmic vesicle, secretory vesicle membrane ; Peripheral membrane protein . Mitochondrion membrane; Peripheral membrane protein.
Purification	The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen.
Clonality	Polyclonal
Concentration	1 mg/ml
Observed band	71kD
Human Gene ID	26056
Human Swiss-Prot Number	Q9BXF6



Alternative Names

Background

disease:Antibodies against RIP11 are found in sera from patients with systemic lupus erythematosus (SLE) or Sjogren syndrome (SS), and in the sera from mothers of children with neonatal lupus erythematosus (NLE).,domain:Binds to vesicles enriched in neutral phospholipids via its C2 domain. The interaction is favored by Mg(2+) rather than Ca(2+).,function:Rab effector involved in protein trafficking from apical recycling endosomes to the apical plasma membrane.,PTM:Phosphorylated on serine and threonine residues.,similarity:Contains 1 C2 domain.,subunit:Forms an heterooligomeric complex with RAB11FIP4. Binds NAPG and SSA2. Binds RAB11A that has been activated by GTP binding.,tissue specificity:Detected at low levels in heart, brain, placenta, lung, liver, adipocytes, kidney, spleen, skeletal muscle and pancreas.,