

## **PTPA** rabbit pAb

## Cat No.:ES10226

For research use only

## Overview

Product Name	PTPA rabbit pAb
Host species	Rabbit
Applications	WB;ELISA
Species Cross-Reactivity	Human;Mouse
<b>Recommended dilutions</b>	WB 1:500-2000 ELISA 1:5000-20000
Immunogen	Synthesized peptide derived from part region of
	human protein
Specificity	PTPA Polyclonal Antibody detects endogenous levels
	of protein.
Formulation	Liquid in PBS containing 50% glycerol, 0.5% BSA and
	0.02% sodium azide.
Storage	Store at -20°C. Avoid repeated freeze-thaw cycles.
Protein Name	Serine/threonine-protein phosphatase 2A activator
	(EC 5.2.1.8) (PP2A, subunit B', PR53 isoform)
	(Phosphotyrosyl phosphatase activator) (PTPA)
	(Serine/threonine-protein phosphatase 2A
	regulatory subuni
Gene Name	PPP2R4 PTPA
Cellular localization	Cytoplasm . Nucleus .
Purification	The antibody was affinity-purified from rabbit
	antiserum by affinity-chromatography using
	epitope-specific immunogen.
Clonality	Polyclonal
Concentration	1 mg/ml
Observed band	39kD
Human Gene ID	5524
Human Swiss-Prot Number	Q15257
Alternative Names	
Background	Protein phosphatase 2A is one of the four major
	Ser/Thr phosphatases and is implicated in the
	negative control of cell growth and division. Protein
	phosphatase 2A holoenzymes are heterotrimeric
	proteins composed of a structural subunit A, a



+86-27-59760950

ELKbio@ELKbiotech.com


www.elkbiotech.com

23-2, No.388 Gaoxin 2nd Road, Wuhan East Lake Hi-tech Development Zone, Hubei , P.R.C



catalytic subunit C, and a regulatory subunit B. The regulatory subunit is encoded by a diverse set of genes that have been grouped into the B/PR55, B'/PR61, and B''/PR72 families. These different regulatory subunits confer distinct enzymatic specificities and intracellular localizations to the holozenzyme. The product of this gene belongs to the B' family. This gene encodes a specific phosphotyrosyl phosphatase activator of the dimeric form of protein phosphatase 2A. Alternative splicing results in multiple transcript variants encoding different isoforms. [provided by RefSeq, Jul 2008],

Western blot analysis of lysates from SW480 cells, primary antibody was diluted at 1:1000, 4°over night





+86-27-59760950

ELKbio@ELKbiotech.com

www.elkbiotech.com

23-2, No.388 Gaoxin 2nd Road, Wuhan East Lake Hi-tech Development Zone, Hubei , P.R.C