

## TAF7 rabbit pAb

Cat No.: ES10382

For research use only

## Overview

Product Name TAF7 rabbit pAb

Host species Rabbit
Applications WB;ELISA
Species Cross-Reactivity Human;Mouse

Recommended dilutions WB 1:500-2000 ELISA 1:5000-20000

**Immunogen** Synthesized peptide derived from part region of

human protein

**Specificity** TAF7 Polyclonal Antibody detects endogenous levels

of protein.

Formulation Liquid in PBS containing 50% glycerol, 0.5% BSA and

0.02% sodium azide.

Storage Store at -20°C. Avoid repeated freeze-thaw cycles.

Protein Name Transcription initiation factor TFIID subunit 7 (RNA polymerase II TBP-associated factor subunit F)

(Transcription initiation factor TFIID 55 kDa subunit)

(TAF(II)55) (TAFII-55) (TAFII55)

Gene Name TAF7 TAF2F TAFII55

Cellular localization Nucleus.

**Purification** The antibody was affinity-purified from rabbit

antiserum by affinity-chromatography using

epitope-specific immunogen.

Clonality Polyclonal
Concentration 1 mg/ml
Observed band 38kD
Human Gene ID 6879
Human Swiss-Prot Number Q15545

**Alternative Names** 

**Background** The intronless gene for this transcription coactivator

is located between the protocadherin beta and gamma gene clusters on chromosome 5. The protein encoded by this gene is a component of the TFIID protein complex, a complex which binds to the TATA

box in class II promoters and recruits RNA




+86-27-59760950

ELKbio@ELKbiotech.com

www.elkbiotech.com





polymerase II and other factors. This particular subunit interacts with the largest TFIID subunit, as well as multiple transcription activators. The protein is required for transcription by promoters targeted by RNA polymerase II. [provided by RefSeq, Jul 2008],

Western blot analysis of lysates from K562 cells, primary antibody was diluted at 1:1000, 4° over night



+86-27-59760950