

HIPK3 rabbit pAb

Cat No.:ES10650

For research use only

Overview

| Product Name | HIPK3 rabbit pAb |
|--------------------------|--|
| Host species | Rabbit |
| Applications | WB;ELISA |
| Species Cross-Reactivity | Human;Rat;Mouse; |
| Recommended dilutions | WB 1:500-2000 ELISA 1:5000-20000 |
| Immunogen | Synthesized peptide derived from human protein . at |
| | AA range: 770-850 |
| Specificity | HIPK3 Polyclonal Antibody detects endogenous |
| | levels of protein. |
| Formulation | Liquid in PBS containing 50% glycerol, 0.5% BSA and |
| | 0.02% sodium azide. |
| Storage | Store at -20 $^\circ\!\mathrm{C}$. Avoid repeated freeze-thaw cycles. |
| Protein Name | Homeodomain-interacting protein kinase 3 (EC |
| | 2.7.11.1) (Androgen receptor-interacting nuclear |
| | protein kinase) (ANPK) (Fas-interacting |
| | serine/threonine-protein kinase) (FIST) (Homolog of |
| | protein kinase |
| Gene Name | HIPK3 DYRK6 FIST3 PKY |
| Cellular localization | Cytoplasm . Nucleus . |
| Purification | The antibody was affinity-purified from rabbit |
| | antiserum by affinity-chromatography using |
| | epitope-specific immunogen. |
| Clonality | Polyclonal |
| Concentration | 1 mg/ml |
| Observed band | 133kD |
| Human Gene ID | 10114 |
| Human Swiss-Prot Number | Q9H422 |
| Alternative Names | |
| Background | catalytic activity:ATP + a protein = ADP + a |
| | phosphoprotein.,function:Seems to negatively |
| | regulate apoptosis by promoting FADD |
| | phosphorylation. Enhances androgen |
| | receptor-mediated transcription. May act as a |
| | |



+86-27-59760950

ELKbio@ELKbiotech.com

www.elkbiotech.com

23-2, No.388 Gaoxin 2nd Road, Wuhan East Lake Hi-tech Development Zone, Hubei , P.R.C



transcriptional corepressor for NK homeodomain transcription factors., PTM: Autophosphorylated, but autophosphorylation is not required for catalytic activity., PTM: May be sumoylated., similarity: Belongs to the protein kinase superfamily. CMGC Ser/Thr protein kinase family. HIPK subfamily., similarity: Contains 1 protein kinase domain., subunit: Interacts with Nkx1-2. Interacts with FAS and DAXX. Probably part of a complex consisting of HIPK3, FAS and FADD. Interacts with and stabilizes ligand-bound androgen receptor (AR) (By similarity). Interacts with UBL1/SUMO-1., tissue specificity: Overexpressed in multidrug resistant cells. Highly expressed in heart and skeletal muscle, and at lower levels in placenta, pancreas, brain, spleen, prostate, thymus, testis, small intestine, colon and leukocytes. Not found in liver and lung.,



+86-27-59760950

ELKbio@ELKbiotech.com

www.elkbiotech.com

23-2, No.388 Gaoxin 2nd Road, Wuhan East Lake Hi-tech Development Zone, Hubei , P.R.C