

SETB2 rabbit pAb

Cat No.:ES10736

For research use only

Overview

Product Name	SETB2 rabbit pAb
Host species	Rabbit
Applications	WB;ELISA
Species Cross-Reactivity	Human;Rat;Mouse;
Recommended dilutions	WB 1:500-2000 ELISA 1:5000-20000
Immunogen	Synthesized peptide derived from part region of
	human protein
Specificity	SETB2 Polyclonal Antibody detects endogenous
	levels of protein.
Formulation	Liquid in PBS containing 50% glycerol, 0.5% BSA and
	0.02% sodium azide.
Storage	Store at -20 $^\circ\!\mathrm{C}$. Avoid repeated freeze-thaw cycles.
Protein Name	Histone-lysine N-methyltransferase SETDB2 (EC
	2.1.1.43) (Chronic lymphocytic leukemia deletion
	region gene 8 protein) (Lysine N-methyltransferase
	1F) (SET domain bifurcated 2)
Gene Name	SETDB2 C13orf4 CLLD8 KMT1F
Cellular localization	Nucleus . Chromosome .
Purification	The antibody was affinity-purified from rabbit
	antiserum by affinity-chromatography using
	epitope-specific immunogen.
Clonality	Polyclonal
Concentration	1 mg/ml
Observed band	79kD
Human Gene ID	83852
Human Swiss-Prot Number	Q96T68
Alternative Names	
Background	SET domain bifurcated 2(SETDB2) Homo sapiens
	This gene encodes a member of a family of proteins
	that contain a methyl-CpG-binding domain (MBD)
	and a SET domain and function as histone
	methyltransferases. This protein is recruited to
	heterochromatin and plays a role in the regulation of



+86-27-59760950

ELKbio@ELKbiotech.com

www.elkbiotech.com

23-2, No.388 Gaoxin 2nd Road, Wuhan East Lake Hi-tech Development Zone, Hubei , P.R.C



chromosome segregation. This region is commonly deleted in chronic lymphocytic leukemia. Naturally-occuring readthrough transcription occurs from this gene to the downstream PHF11 (PHD finger protein 11) gene. Alternative splicing results in multiple transcript variants. [provided by RefSeq, Mar 2016],



+86-27-59760950

ELKbio@ELKbiotech.com

www.elkbiotech.com

23-2, No.388 Gaoxin 2nd Road,Wuhan East Lake Hi-tech Development Zone, Hubei , P.R.C