

## PKN3 rabbit pAb

## Cat No.:ES10826

For research use only

## Overview

| Product Name                 | PKN3 rabbit pAb  |
|------------------------------|--|
| Host species                 | Rabbit   |
| Applications                 | WB;ELISA   |
| Species Cross-Reactivity     | Human;Mouse  |
| Recommended dilutions        | WB 1:500-2000 ELISA 1:5000-20000                                       |
| Immunogen                    | Synthesized peptide derived from part region of                        |
|                              | human protein  |
| Specificity                  | PKN3 Polyclonal Antibody detects endogenous levels                     |
|                              | of protein.  |
| Formulation                  | Liquid in PBS containing 50% glycerol, 0.5% BSA and                    |
|                              | 0.02% sodium azide.  |
| Storage                      | Store at -20 $^\circ\!\mathrm{C}$ . Avoid repeated freeze-thaw cycles. |
| Protein Name                 | Serine/threonine-protein kinase N3 (EC 2.7.11.13)                      |
|                              | (Protein kinase PKN-beta) (Protein-kinase C-related                    |
|                              | kinase 3)  |
| Gene Name                    | PKN3 PKNBETA   |
| <b>Cellular localization</b> | Nucleus . Cytoplasm, perinuclear region . Nuclear                      |
|                              | and perinuclear Golgi region.  |
| Purification                 | The antibody was affinity-purified from rabbit                         |
|                              | antiserum by affinity-chromatography using                             |
|                              | epitope-specific immunogen.  |
| Clonality                    | Polyclonal   |
| Concentration                | 1 mg/ml  |
| Observed band                | 97kD   |
| Human Gene ID                | 29941  |
| Human Swiss-Prot Number      | Q6P5Z2   |
| Alternative Names            |  |
| Background                   | catalytic activity:ATP + a protein = ADP + a                           |
|                              | phosphoprotein.,domain:The C1 domain does not                          |
|                              | bind the diacylglycerol (DAG).,enzyme                                  |
|                              | regulation:Two specific sites, Thr-718 (activation                     |
|                              | loop of the kinase domain) and Thr-860 (turn motif),                   |
|                              | need to be phosphorylated for its full                                 |



+86-27-59760950

ELKbio@ELKbiotech.com

www.elkbiotech.com

23-2, No.388 Gaoxin 2nd Road, Wuhan East Lake Hi-tech Development Zone, Hubei , P.R.C



activation.,function:Contributes to invasiveness in malignant prostate cancer.,PTM:Autophosphorylated.,similarity:Belongs to the protein kinase superfamily. AGC Ser/Thr protein kinase family. PKC subfamily.,similarity:Contains 1 AGC-kinase C-terminal domain.,similarity:Contains 1 protein kinase domain.,similarity:Contains 3 REM (Hr1) repeats.,subcellular location:Nuclear and perinuclear Golgi region.,tissue specificity:Expressed in prostate tumors and various cancer cell lines. Not expressed in adult tissues.,



+86-27-59760950

ELKbio@ELKbiotech.com

www.elkbiotech.com

23-2, No.388 Gaoxin 2nd Road, Wuhan East Lake Hi-tech Development Zone, Hubei , P.R.C