

UVRAG rabbit pAb

Cat No.:ES10871

For research use only

Overview

| Product Name | UVRAG rabbit pAb |
|------------------------------|--|
| Host species | Rabbit |
| Applications | WB;ELISA |
| Species Cross-Reactivity | Human;Rat;Mouse; |
| Recommended dilutions | WB 1:500-2000 ELISA 1:5000-20000 |
| Immunogen | Synthesized peptide derived from part region of |
| | human protein. AA range 655-675 |
| Specificity | UVRAG Polyclonal Antibody detects endogenous |
| | levels of protein. |
| Formulation | Liquid in PBS containing 50% glycerol, 0.5% BSA and |
| | 0.02% sodium azide. |
| Storage | Store at -20 $^\circ\!{ m C}$. Avoid repeated freeze-thaw cycles. |
| Protein Name | UV radiation resistance-associated gene protein |
| | (p63) |
| Gene Name | UVRAG |
| Cellular localization | Late endosome . Lysosome . Cytoplasmic vesicle, |
| | autophagosome . Early endosome . Endoplasmic |
| | reticulum . Midbody . Chromosome, centromere . |
| | Colocalizes with RAB9-positive compartments |
| | involved in retrograde transport from late |
| | endosomes to trans-Golgi network. Colocalization |
| | with early endosomes is only partial |
| | (PubMed:24056303). Recruited to autophagosome |
| | following interaction with RUBCNL/PACER |
| | (PubMed:28306502) |
| Purification | The antibody was affinity-purified from rabbit |
| | antiserum by affinity-chromatography using |
| | epitope-specific immunogen. |
| Clonality | Polyclonal |
| Concentration | 1 mg/ml |
| Observed band | 76kD |
| Human Gene ID | 7405 |
| Human Swiss-Prot Number | Q9P2Y5 |
| | |



+86-27-59760950

ELKbio@ELKbiotech.com

www.elkbiotech.com

23-2, No.388 Gaoxin 2nd Road, Wuhan East Lake Hi-tech Development Zone, Hubei , P.R.C



Alternative Names Background

This gene complements the ultraviolet sensitivity of xeroderma pigmentosum group C cells and encodes a protein with a C2 domain. The protein activates the Beclin1-PI(3)KC3 complex, promoting autophagy and suppressing the proliferation and tumorigenicity of human colon cancer cells. Chromosomal aberrations involving this gene are associated with left-right axis malformation and mutations in this gene have been associated with colon cancer. [provided by RefSeq, Jul 2008],



+86-27-59760950

ELKbio@ELKbiotech.com

www.elkbiotech.com

23-2, No.388 Gaoxin 2nd Road, Wuhan East Lake Hi-tech Development Zone, Hubei , P.R.C