

CO5 rabbit pAb

Cat No.:ES11173

For research use only

Overview

Product Name	CO5 rabbit pAb
Host species	Rabbit
Applications	WB;ELISA
Species Cross-Reactivity	Human;Rat;Mouse
Recommended dilutions	WB 1:500-2000 ELISA 1:5000-20000
Immunogen	Synthesized peptide derived from human protein . at
	AA range: 690-770
Specificity	CO5 Polyclonal Antibody detects endogenous levels
	of protein.
Formulation	Liquid in PBS containing 50% glycerol, 0.5% BSA and
	0.02% sodium azide.
Storage	Store at -20 $^\circ\!\mathbb{C}$. Avoid repeated freeze-thaw cycles.
Protein Name	Complement C5 (C3 and PZP-like
	alpha-2-macroglobulin domain-containing protein 4)
	[Cleaved into: Complement C5 beta chain;
	Complement C5 alpha chain; C5a anaphylatoxin;
	Complement C5 alpha' chain]
Gene Name	C5 CPAMD4
Cellular localization	Secreted.
Purification	The antibody was affinity-purified from rabbit
	antiserum by affinity-chromatography using
	epitope-specific immunogen.
Clonality	Polyclonal
Concentration	1 mg/ml
Observed band	184kD
Human Gene ID	727
Human Swiss-Prot Number	P01031
Alternative Names	
Background	This gene encodes a component of the complement
	system, a part of the innate immune system that
	plays an important role in inflammation, host
	homeostasis, and host defense against pathogens.
	The encoded preproprotein is proteolytically



+86-27-59760950

ELKbio@ELKbiotech.com

www.elkbiotech.com

23-2, No.388 Gaoxin 2nd Road, Wuhan East Lake Hi-tech Development Zone, Hubei , P.R.C



processed to generate multiple protein products, including the C5 alpha chain, C5 beta chain, C5a anaphylatoxin and C5b. The C5 protein is comprised of the C5 alpha and beta chains, which are linked by a disulfide bridge. Cleavage of the alpha chain by a convertase enzyme results in the formation of the C5a anaphylatoxin, which possesses potent spasmogenic and chemotactic activity, and the C5b macromolecular cleavage product, a subunit of the membrane attack complex (MAC). Mutations in this gene cause complement component 5 deficiency, a disease characterized by recurrent bacterial infections. Alternative splicing results in multiple transcript variants. [provided by RefSeq,



+86-27-59760950

ELKbio@ELKbiotech.com

www.elkbiotech.com

23-2, No.388 Gaoxin 2nd Road, Wuhan East Lake Hi-tech Development Zone, Hubei , P.R.C