



# TNKS2 rabbit pAb

Cat No.:ES11342

For research use only

## Overview

Product Name	TNKS2 rabbit pAb
Host species	Rabbit
Applications	WB;ELISA
Species Cross-Reactivity	Human;Mouse
Recommended dilutions	WB 1:500-2000 ELISA 1:5000-20000
Immunogen	Synthesized peptide derived from human protein . at AA range: 950-1030
Specificity	TNKS2 Polyclonal Antibody detects endogenous levels of protein.
Formulation	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.
Storage	Store at -20℃ . Avoid repeated freeze-thaw cycles.
Protein Name	Tankyrase-2 (TANK2) (EC 2.4.2.30) (ADP-ribosyltransferase diphtheria toxin-like 6) (ARTD6) (Poly [ADP-ribose] polymerase 5B) (TNKS-2) (TRF1-interacting ankyrin-related ADP-ribose polymerase 2) (Tankyr
Gene Name	TNKS2 PARP5B TANK2 TNKL
Cellular localization	Cytoplasm. Golgi apparatus membrane; Peripheral membrane protein. Nucleus. Chromosome, telomere . Associated with the Golgi and with juxtannuclear SLC2A4/GLUT4-vesicles. Also found around the pericentriolar matrix of mitotic centromeres. During interphase, a small fraction of TNKS2 is found in the nucleus, associated with TRF1.
Purification	The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen.
Clonality	Polyclonal
Concentration	1 mg/ml
Observed band	128kD
Human Gene ID	80351
Human Swiss-Prot Number	Q9H2K2





## Alternative Names Background

catalytic activity:NAD(+) + (ADP-D-ribosyl)(n)-acceptor = nicotinamide + (ADP-D-ribosyl)(n+1)-acceptor.,function:May regulate vesicle trafficking and modulate the subcellular distribution of SLC2A4/GLUT4-vesicles. Has PARP activity and can modify TRF1, and thereby contribute to the regulation of telomere length.,PTM:ADP-ribosylated (-auto).,similarity:Contains 1 PARP catalytic domain.,similarity:Contains 1 SAM (sterile alpha motif) domain.,similarity:Contains 15 ANK repeats.,subcellular location:Associated with the Golgi and with juxtannuclear SLC2A4/GLUT4-vesicles. Also found around the pericentriolar matrix of mitotic centromeres. During interphase, a small fraction of TNKS2 is found in the nucleus, associated with TRF1.,subunit:Oligomerizes and associates with TNKS. Interacts with the cytoplasmic domain of LNPEP/Otase in SLC2A4/GLUT4-vesicles. Binds to the N-terminus of Grb14 and TRF1 with its ankyrin repeat region.,tissue specificity:Highly expressed in placenta, skeletal muscle, liver, brain, kidney, heart, thymus, spinal cord, lung, peripheral blood leukocytes, pancreas, lymph nodes, spleen, prostate, testis, ovary, small intestine, colon, mammary gland, breast and breast carcinoma, and in common-type meningioma. Highly expressed in fetal liver, heart and brain.,

